

thm_2Esat_2Epth__no2
(TMQu8ewnnFJhVj62ELjGs6GX8CxyDJK62kv)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 6 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F V0t))$

Definition 8 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \vee (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \vee True) \Leftrightarrow True) \wedge \\ & (((False \vee (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee False) \Leftrightarrow (p V0t)) \wedge (((p V0t) \vee \\ & (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \tag{4}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (5)$$

Theorem 1 $(\forall V0p \in 2.(\forall V1q \in 2.(\neg((p V0p) \vee (p V1q))) \Rightarrow (\neg(p V1q))))$.