

# thm\_2Esemi\_ring\_2Eplus\_rotate (TMUb- NiSuMLPKRqEb6wqUnwEBUM6keKwRfh2)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Ebool_2E_2T` to be  $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let `ty_2Esemi_ring_2Esemi_ring` :  $\iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Esemi\_ring\_2Esemi\_ring A0) \quad (1)$$

Let `c_2Esemi_ring_2Esemi_ring_SRM` :  $\iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow c\_2Esemi\_ring\_2Esemi\_ring\_SRM A.27a \in (A.27a^{(ty\_2Esemi\_ring\_2Esemi\_ring A.27a)}) \quad (2)$$

Let `c_2Esemi_ring_2Esemi_ring_SRM0` :  $\iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow c\_2Esemi\_ring\_2Esemi\_ring\_SRM0 A.27a \in (A.27a^{(ty\_2Esemi\_ring\_2Esemi\_ring A.27a)}) \quad (3)$$

Let `c_2Esemi_ring_2Esemi_ring_SRM` :  $\iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow c\_2Esemi\_ring\_2Esemi\_ring\_SRM A.27a \in (((A.27a^{A.27a})^{A.27a})^{(ty\_2Esemi\_ring\_2Esemi\_ring A.27a)}) \quad (4)$$

Let `c_2Esemi_ring_2Esemi_ring_SRP` :  $\iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow c\_2Esemi\_ring\_2Esemi\_ring\_SRP A.27a \in (((A.27a^{A.27a})^{A.27a})^{(ty\_2Esemi\_ring\_2Esemi\_ring A.27a)}) \quad (5)$$

**Definition 4** We define `c_2Ebool_2E_21` to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A.27a}).(ap (ap (c_2Emin_2E_3D (2^{A.27a}))$

**Definition 5** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.$

**Definition 6** We define  $c\_2Esemi\_ring\_2Eis\_semi\_ring$  to be  $\lambda A\_27a : \iota.\lambda V0r \in (ty\_2Esemi\_ring\_2Esemi\_ring$

Assume the following.

$$True \tag{6}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{7}$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0r \in (ty\_2Esemi\_ring\_2Esemi\_ring \\ & \quad A\_27a). ((p\ (ap\ (c\_2Esemi\_ring\_2Eis\_semi\_ring\ A\_27a)\ V0r)) \Rightarrow \\ & \quad (\forall V1m \in A\_27a. (\forall V2n \in A\_27a. (\forall V3p \in A\_27a. ( \\ & \quad (ap\ (ap\ (ap\ (c\_2Esemi\_ring\_2Esemi\_ring\_SRP\ A\_27a)\ V0r)\ (ap \\ & \quad (ap\ (ap\ (c\_2Esemi\_ring\_2Esemi\_ring\_SRP\ A\_27a)\ V0r)\ V1m)\ V2n)) \\ & \quad V3p) = (ap\ (ap\ (ap\ (c\_2Esemi\_ring\_2Esemi\_ring\_SRP\ A\_27a)\ V0r) \\ & \quad (ap\ (ap\ (ap\ (c\_2Esemi\_ring\_2Esemi\_ring\_SRP\ A\_27a)\ V0r)\ V2n) \\ & \quad V3p))\ V1m)))))) \end{aligned}$$