

thm\_2Eseq\_2EK\_PARTIAL (TM-  
coxD5k3TDwPc6WStWudpssB7sA1KhkHAY)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ecombin\_2EK$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.(\lambda V0x \in A\_27a.(\lambda V1y \in A\_27b.V0x))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \tag{2}$$

**Theorem 1**

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow (\forall V0x \in A\_27a.((ap (c\_2Ecombin\_2EK A\_27a A\_27b) V0x) = (\lambda V1z \in A\_27b.V0x)))$$