

thm_2Esorting_2EPERM__EQUIVALENCE__ALT__DEF
(TMK_wH-
BGF_XV_nP₄R_KW_Mx_{is}H_XH_LT_H1_xE_v37_pe_x)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)$

Let $ty_2Elist_2Elist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Elist_2Elist A0) \quad (1)$$

Let $c_2Elist_2EFILTER : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2EFILTER A_27a \in (((ty_2Elist_2Elist A_27a)(ty_2Elist_2Elist A_27a))(2^{A_27a})) \quad (2)$$

Definition 8 We define $c_2Esorting_2EPERM$ to be $\lambda A_27a : \iota.\lambda V0L1 \in (ty_2Elist_2Elist A_27a).\lambda V1L2 \in (ty_2Elist_2Elist A_27a)$

Definition 9 We define $c_2Erelation_2Etransitive$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_7E$

Definition 10 We define $c_2Erelation_2Esymmetric$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_7E$

Definition 11 We define $c_2Erelation_2Ereflexive$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E$

Definition 12 We define $c_2Erelation_2Eequivalence$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (ap c_2E$

Assume the following.

$$True \quad (3)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (4)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\ & p V0t)))))) \quad (5) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0R \in ((2^{A_27a})^{A_27a}). \\ & ((p (ap (c_2Erelation_2Eequivalence A_27a) V0R)) \Leftrightarrow (\forall V1x \in \\ & A_27a. (\forall V2y \in A_27a. ((p (ap (ap V0R V1x) V2y)) \Leftrightarrow ((ap V0R V1x) = \\ & (ap V0R V2y))))))) \quad (6) \end{aligned}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (p (ap (c_2Erelation_2Eequivalence (ty_2Elist_2Elist A_27a) (c_2Esorting_2Eperm A_27a))) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in (ty_2Elist_2Elist \\ & A_27a). (\forall V1y \in (ty_2Elist_2Elist A_27a). ((p (ap (ap (c_2Esorting_2Eperm \\ & A_27a) V0x) V1y)) \Leftrightarrow ((ap (c_2Esorting_2Eperm A_27a) V0x) = (ap (c_2Esorting_2Eperm \\ & A_27a) V1y)))))) \end{aligned}$$