

thm_2Estring_2ECHAR__EQ__THM
(TMNcHwffC41oYiw7oBGBHP1d1uFVyrD8f7Y)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $ty_2Estring_2Echar : \iota$ be given. Assume the following.

$$nonempty\ ty_2Estring_2Echar \tag{2}$$

Let $c_2Estring_2EORD : \iota$ be given. Assume the following.

$$c_2Estring_2EORD \in (ty_2Enum_2Enum^{ty_2Estring_2Echar}) \tag{3}$$

Assume the following.

$$True \tag{4}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \tag{5}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (6)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (7)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (8)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (9)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_27 \in 2.(\forall V2y \in 2.(\forall V3y_27 \in 2.(((p V0x) \Leftrightarrow (p V1x_27)) \wedge ((p V1x_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_27)))))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_27) \Rightarrow (p V3y_27)))))) \quad (10)$$

Assume the following.

$$(\forall V0a \in ty_2Estring_2Echar.(\forall V1a_27 \in ty_2Estring_2Echar.(((ap c_2Estring_2EORD V0a) = (ap c_2Estring_2EORD V1a_27)) \Leftrightarrow (V0a = V1a_27)))) \quad (11)$$

Theorem 1

$$(\forall V0c1 \in ty_2Estring_2Echar.(\forall V1c2 \in ty_2Estring_2Echar.(((V0c1 = V1c2) \Leftrightarrow ((ap c_2Estring_2EORD V0c1) = (ap c_2Estring_2EORD V1c2))))))$$