

thm_2Estring_2EORD__11 (TMKge- BuXLSvwkh1gcjpMUnAqnMqNponwDEu)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow p \Rightarrow Q)$ of type ι .

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty \ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Enum_2E0 to be $(ap \ c_2Enum_2EABS_num \ c_2Enum_2EZERO_REP)$.

Definition 4 We define $c_2Earithmic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 5 We define c_2Ebool_2ET to be $(ap \ (ap \ (c_2Emin_2E_3D \ (2^2)) \ (\lambda V0x \in 2.V0x)) \ (\lambda V1x \in 2.V1x))$

Definition 6 We define $c_2Ebool_2E_21$ to be $\lambda A.^{27a} : \iota.(\lambda V0P \in (2^{A-27a}).(ap \ (ap \ (c_2Emin_2E_3D \ (2^{A-27a})))$

Definition 7 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap \ c_2Enum_2EABS_num \ (ap \ c_2Enum_2EREP_num \ c_2Enum_2ESUC_REP))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 8 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2E_2B V0n))$.

Definition 9 We define $c_2Earithmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2E_2B V0n))$.

Definition 10 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Definition 11 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 12 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E))$.

Definition 13 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$.

Definition 14 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 15 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a P))))$.

Definition 16 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.V1n$.

Let $ty_2Estring_2Echar : \iota$ be given. Assume the following.

$$nonempty\ ty_2Estring_2Echar \quad (7)$$

Let $c_2Estring_2EORD : \iota$ be given. Assume the following.

$$c_2Estring_2EORD \in (ty_2Enum_2Enum^{ty_2Estring_2Echar}) \quad (8)$$

Let $c_2Estring_2ECHR : \iota$ be given. Assume the following.

$$c_2Estring_2ECHR \in (ty_2Estring_2Echar^{ty_2Enum_2Enum}) \quad (9)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (10)$$

Assume the following.

$$\begin{aligned} & ((\forall V0a \in ty_2Estring_2Echar.((ap c_2Estring_2ECHR (ap \\ & c_2Estring_2EORD V0a) = V0a)) \wedge (\forall V1r \in ty_2Enum_2Enum. \\ & ((p (ap (\lambda V2n \in ty_2Enum_2Enum.(ap (ap c_2Eprim_rec_2E_3C \\ & V2n) (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 \\ & (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 \\ & (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 \\ & (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO)))))))))) \\ & V1r)) \Leftrightarrow ((ap c_2Estring_2EORD (ap c_2Estring_2ECHR V1r) = V1r)))) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} & (\forall V0a \in ty_2Estring_2Echar.(\forall V1a_27 \in ty_2Estring_2Echar. \\ & (((ap c_2Estring_2EORD V0a) = (ap c_2Estring_2EORD V1a_27)) \Leftrightarrow (\\ & V0a = V1a_27)))) \end{aligned}$$