

# thm\_2Estring\_2EORD\_\_CHR (TMXzZa- ehd4xGRADLoq7JnSGvwPMt3Lr83Gy)

October 26, 2020

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

**Definition 3** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

**Definition 4** We define  $c\_2Ebool\_2ET$  to be  $(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

**Definition 5** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

**Definition 6** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num\ (ap\ (ap\ (c\_2Emin\_2E\_3D\ (2^{2^m}))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{6}$$

**Definition 7** We define  $c\_2Earithmic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmic\_2EBIT1 V0n) (ap c\_2Earithmic\_2EBIT1 V0n))$ .

**Definition 8** We define  $c\_2Earithmic\_2EBIT2$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmic\_2EBIT2 V0n) (ap c\_2Earithmic\_2EBIT2 V0n))$ .

**Definition 9** We define  $c\_2Earithmic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

**Definition 10** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 11** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 12** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_7E V0t))$ .

**Definition 13** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t) V1t2) V0t1))$ .

**Definition 14** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 15** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\lambda P \in 2^A.\lambda V0P \in (2^A)^{2^A}.(ap V0P (ap (c\_2Emin\_2E\_40 V0P)))$ .

**Definition 16** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum.V1n$ .

Let  $ty\_2Estring\_2Echar : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Estring\_2Echar \quad (7)$$

Let  $c\_2Estring\_2EORD : \iota$  be given. Assume the following.

$$c\_2Estring\_2EORD \in (ty\_2Enum\_2Enum^{ty\_2Estring\_2Echar}) \quad (8)$$

Let  $c\_2Estring\_2ECHR : \iota$  be given. Assume the following.

$$c\_2Estring\_2ECHR \in (ty\_2Estring\_2Echar^{ty\_2Enum\_2Enum}) \quad (9)$$

Assume the following.

$$\begin{aligned} & ((\forall V0a \in ty\_2Estring\_2Echar.((ap c\_2Estring\_2ECHR (ap \\ & c\_2Estring\_2EORD V0a)) = V0a)) \wedge (\forall V1r \in ty\_2Enum\_2Enum. \\ & ((p (ap (\lambda V2n \in ty\_2Enum\_2Enum.(ap (ap c\_2Eprim\_rec\_2E\_3C \\ & V2n) (ap c\_2Earithmic\_2ENUMERAL (ap c\_2Earithmic\_2EBIT2 \\ & (ap c\_2Earithmic\_2EBIT1 (ap c\_2Earithmic\_2EBIT1 (ap c\_2Earithmic\_2EBIT1 \\ & (ap c\_2Earithmic\_2EBIT1 (ap c\_2Earithmic\_2EBIT1 (ap c\_2Earithmic\_2EBIT1 \\ & (ap c\_2Earithmic\_2EBIT1 c\_2Earithmic\_2EZERO)))))))))) \\ & V1r)) \Leftrightarrow ((ap c\_2Estring\_2EORD (ap c\_2Estring\_2ECHR V1r)) = V1r))) \end{aligned} \quad (10)$$

**Theorem 1**

$$\begin{aligned} & ((\forall V0r \in ty\_2Enum\_2Enum.((p (ap (ap c\_2Eprim\_rec\_2E\_3C \\ & V0r) (ap c\_2Earithmic\_2ENUMERAL (ap c\_2Earithmic\_2EBIT2 \\ & (ap c\_2Earithmic\_2EBIT1 (ap c\_2Earithmic\_2EBIT1 (ap c\_2Earithmic\_2EBIT1 \\ & (ap c\_2Earithmic\_2EBIT1 (ap c\_2Earithmic\_2EBIT1 (ap c\_2Earithmic\_2EBIT1 \\ & (ap c\_2Earithmic\_2EBIT1 c\_2Earithmic\_2EZERO)))))))))) \Leftrightarrow \\ & ((ap c\_2Estring\_2EORD (ap c\_2Estring\_2ECHR V0r)) = V0r))) \end{aligned}$$