

thm\_2Esum\_2EEXISTS\_SUM  
(TMQv4gWSH6Yg3BUaX7LPj2fFcnsGJ3hQ4xL)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  then  $(the (\lambda x.x \in A \wedge p (ap P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A\_27a P))$

**Definition 7** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 8** We define  $c\_2Ebool\_2E\_5C\_2E\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 9** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 10** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Esum\_2Esum A0 A1) \tag{1}$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Esum\_2EABS\_sum A\_27a A\_27b \in ((ty\_2Esum\_2Esum A\_27a A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \tag{2}$$

**Definition 11** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27b.(ap (c\_2Esum\_2EABS\_sum A\_27a A\_27b) V0e)$

**Definition 12** We define  $c\_2Esum\_2EINL$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27a.(ap (c\_2Esum\_2EABS$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (3)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}).((\neg(\forall V1x \in A\_27a.(p (ap V0P V1x)))) \Leftrightarrow (\exists V2x \in A\_27a.(\neg(p (ap V0P V2x)))))) \quad (4)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \wedge (p V1B))) \Leftrightarrow ((\neg(p V0A) \vee \neg(p V1B)))) \wedge (((\neg((p V0A) \vee (p V1B))) \Leftrightarrow ((\neg(p V0A) \wedge \neg(p V1B))))))) \quad (5)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow (\forall V0P \in (2^{(ty\_2Esum\_2Esum A\_27a A\_27b)}).((\forall V1s \in (ty\_2Esum\_2Esum A\_27a A\_27b).(p (ap V0P V1s))) \Leftrightarrow ((\forall V2x \in A\_27a.(p (ap V0P (ap (c\_2Esum\_2EINL A\_27a A\_27b) V2x)))) \wedge (\forall V3y \in A\_27b.(p (ap V0P (ap (c\_2Esum\_2EINR A\_27a A\_27b) V3y)))))))) \quad (6)$$

**Theorem 1**

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow (\forall V0P \in (2^{(ty\_2Esum\_2Esum A\_27a A\_27b)}).((\exists V1s \in (ty\_2Esum\_2Esum A\_27a A\_27b).(p (ap V0P V1s))) \Leftrightarrow ((\exists V2x \in A\_27a.(p (ap V0P (ap (c\_2Esum\_2EINL A\_27a A\_27b) V2x)))) \vee (\exists V3y \in A\_27b.(p (ap V0P (ap (c\_2Esum\_2EINR A\_27a A\_27b) V3y))))))))$$