

thm_2Esum_2EINL_11 (TMKGICYXD- DEZv3pUqEc7T69L6U2kAs9n3put)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \ P \Rightarrow p \ Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

Definition 5 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 7 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.(ap (c_2Ebool_2E_7E V2t) c_2Ebool_2EF))))$

Definition 8 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (\text{the } (\lambda x.x \in A \wedge p x)) \text{ else } (\lambda x.x \in A \wedge \neg p x)$ of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a) (\lambda V1P \in 2.V1P))))$

Definition 10 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.(ap (c_2Ebool_2E_3F V2t) c_2Ebool_2EF))))$

Definition 11 We define $c_2Esum_2EIS_SUM_REP$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (((2^{A_27b})^{A_27a})^2).$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall A0.\text{nonempty } A0 \Rightarrow \forall A1.\text{nonempty } A1 \Rightarrow \text{nonempty } (ty_2Esum_2Esum \\ & \quad A0 \ A1) \end{aligned} \tag{1}$$

Let $c_2Esum_2EREPP_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow c_2Esum_2EREPP_sum \\ & \quad A_27a \ A_27b \in (((2^{A_27b})^{A_27a})^2)^{(ty_2Esum_2Esum \ A_27a \ A_27b)}) \end{aligned} \tag{2}$$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow c_2Esum_2EABS_sum \\ A_27a \ A_27b \in ((ty_2Esum_2Esum\ A_27a\ A_27b)^{((2^{A_27b})^{A_27a})^2}) \end{aligned} \quad (3)$$

Definition 12 We define c_2Esum_2EINL to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27a. (ap (c_2Esum_2EABS_sum$

Assume the following.

$$True \quad (4)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0t \in 2. ((\exists V1x \in \\ A_27a. (p\ V0t) \Leftrightarrow (p\ V0t))) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow \\ True)) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (10)$$

Assume the following.

$$\begin{aligned} \forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow \\ (\forall V0a \in (ty_2Esum_2Esum\ A_27a\ A_27b). ((ap (c_2Esum_2EABS_sum \\ A_27a\ A_27b) (ap (c_2Esum_2EREP_sum\ A_27a\ A_27b)\ V0a)) = V0a)) \wedge \\ (\forall V1r \in ((2^{A_27b})^{A_27a})^2). ((p (ap (c_2Esum_2EIS_SUM_REP \\ A_27a\ A_27b)\ V1r)) \Leftrightarrow ((ap (c_2Esum_2EREP_sum\ A_27a\ A_27b) (ap (\\ c_2Esum_2EABS_sum\ A_27a\ A_27b)\ V1r)) = V1r))) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow (\\ & \quad \forall V0x \in A_27a.(\forall V1y \in A_27a.(((ap(c_2Esum_2EINL A_27a \\ & \quad A_27b) V0x) = (ap(c_2Esum_2EINL A_27a A_27b) V1y)) \Leftrightarrow (V0x = V1y))) \end{aligned}$$