

# thm\_2Esum\_2EINR

## (TMU4PfdsZEZSdEpJf11StkUvVTppHczBFfQ)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A P))$

**Definition 8** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Esum\_2Esum A0 A1) \tag{1}$$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Esum\_2EABS\_sum A\_27a A\_27b \in ((ty\_2Esum\_2Esum A\_27a A\_27b)^{((2^{A-27b})^{A-27a})^2}) \tag{2}$$

**Definition 10** We define  $c\_2Esum\_2EINL$  to be  $\lambda A.\lambda a : \iota.\lambda A.\lambda b : \iota.\lambda V0e \in A.\lambda a.(ap (c\_2Esum\_2EABS\_sum A a) V0e)$

**Definition 11** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_5C\_2F V0t))$

Let  $c\_2Esum\_2EISR : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EISR \\ A\_27a\ A\_27b \in (2^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)}) \end{aligned} \quad (3)$$

**Definition 12** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0e \in A\_27b. (ap\ (c\_2Esum\_2EABS$

Let  $c\_2Esum\_2EOUTR : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EOUTR \\ A\_27a\ A\_27b \in (A\_27b^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)}) \end{aligned} \quad (4)$$

Assume the following.

$$True \quad (5)$$

Assume the following.

$$\begin{aligned} (\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \end{aligned} \quad (6)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p\ V0t))) \quad (7)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (9)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg( \\ p\ V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ \forall V0ss \in (ty\_2Esum\_2Esum\ A\_27a\ A\_27b). ((\exists V1x \in A\_27a. \\ (V0ss = (ap\ (c\_2Esum\_2EINL\ A\_27a\ A\_27b)\ V1x))) \vee (\exists V2y \in A\_27b. \\ (V0ss = (ap\ (c\_2Esum\_2EINR\ A\_27a\ A\_27b)\ V2y)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ & (\forall V0x \in A\_27b.(p\ (ap\ (c\_2Esum\_2EISR\ A\_27a\ A\_27b)\ (ap\ (c\_2Esum\_2EINR \\ & A\_27a\ A\_27b)\ V0x)))) \wedge (\forall V1y \in A\_27a.(\neg(p\ (ap\ (c\_2Esum\_2EISR \\ & A\_27a\ A\_27b)\ (ap\ (c\_2Esum\_2EINL\ A\_27a\ A\_27b)\ V1y)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ & \forall V0x \in A\_27b.((ap\ (c\_2Esum\_2EOUTR\ A\_27a\ A\_27b)\ (ap\ (c\_2Esum\_2EINR \\ & A\_27a\ A\_27b)\ V0x)) = V0x)) \end{aligned} \quad (13)$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ & \forall V0x \in (ty\_2Esum\_2Esum\ A\_27a\ A\_27b).((p\ (ap\ (c\_2Esum\_2EISR \\ & A\_27a\ A\_27b)\ V0x)) \Rightarrow ((ap\ (c\_2Esum\_2EINR\ A\_27a\ A\_27b)\ (ap\ (c\_2Esum\_2EOUTR \\ & A\_27a\ A\_27b)\ V0x)) = V0x))) \end{aligned}$$