

thm_2Esum_2EINR_neq_INL (TMSgaUDm- mJqAj1fiaSuAjHRTnRYXJHWVKLn)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$.

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 7 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then}$ (the $(\lambda x.x \in A \wedge p (ap P x))$ of type $\iota \Rightarrow \iota$.

Definition 8 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a P)))$

Definition 9 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 10 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Definition 11 We define $c_2Esum_2EIS_SUM_REP$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (((2^{A_27b})^{A_27a})^2)$.

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Esum_2Esum A0 A1) \tag{1}$$

Let $c_2Esum_2EREP_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Esum_2EREP_sum A_27a A_27b \in (((2^{A_27b})^{A_27a})^2)^{(ty_2Esum_2Esum A_27a A_27b)} \tag{2}$$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EABS_sum \\ A_27a\ A_27b \in ((ty_2Esum_2Esum\ A_27a\ A_27b)^{((2^{A_27b})^{A_27a})^2}) \end{aligned} \quad (3)$$

Definition 12 We define c_2Esum_2EINL to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27a.(ap\ (c_2Esum_2EABS_sum\ V0e)\ A_27b)$

Definition 13 We define c_2Esum_2EINR to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27b.(ap\ (c_2Esum_2EABS_sum\ V0e)\ A_27a)$

Assume the following.

$$True \quad (4)$$

Assume the following.

$$\begin{aligned} (\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \end{aligned} \quad (5)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (6)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\exists V1x \in \\ A_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} ((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge \\ ((\neg False) \Leftrightarrow True))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow \\ True)) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(\\
& p V0t))))))
\end{aligned} \tag{12}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\
& (\forall V0a \in (ty_2Esum_2Esum A_27a A_27b).((ap (c_2Esum_2EABS_sum \\
& A_27a A_27b) (ap (c_2Esum_2EREP_sum A_27a A_27b) V0a)) = V0a)) \wedge \\
& (\forall V1r \in (((2^{A_27b})^{A_27a})^2).((p (ap (c_2Esum_2EIS_SUM_REP \\
& A_27a A_27b) V1r)) \Leftrightarrow ((ap (c_2Esum_2EREP_sum A_27a A_27b) (ap (\\
& c_2Esum_2EABS_sum A_27a A_27b) V1r)) = V1r))))
\end{aligned} \tag{13}$$

Theorem 1

$$\begin{aligned}
& \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\
& \forall V0v1 \in A_27a.(\forall V1v2 \in A_27b.(\neg((ap (c_2Esum_2EINR \\
& A_27a A_27b) V1v2) = (ap (c_2Esum_2EINL A_27a A_27b) V0v1))))))
\end{aligned}$$