

thm\_2Esum\_2Esum\_\_Axiom  
(TMFi4rfGPQ7NMLAv6rNxxJLKCnB36TBit08)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (P \Rightarrow Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p (ap P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A\_27a P))$

**Definition 8** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Esum\_2Esum A0 A1) \tag{1}$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Esum\_2EABS\_sum A\_27a A\_27b \in ((ty\_2Esum\_2Esum A\_27a A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \tag{2}$$

**Definition 10** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27b.(ap (c\_2Esum\_2EABS\_sum A\_27a A\_27b) V0e)$

**Definition 11** We define  $c\_2Esum\_2EINL$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27a.(ap (c\_2Esum\_2EABS\_sum A\_27a A\_27b) V0e)$

**Definition 12** We define  $c\_2Ecombin\_2Eo$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda A\_27c : \iota.\lambda V0f \in (A\_27b^{A\_27c}).\lambda V1g \in (A\_27c^{A\_27a}).\lambda V2h \in (A\_27a^{A\_27b}).(ap (ap (ap (c\_2Ecombin\_2Eo A\_27c A\_27b A\_27a) V0f) V1g) V2h) = (ap V0f (ap V1g V2h))$

**Definition 13** We define  $c\_2Ebool\_2E\_3F\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap c\_2Ebool\_2E\_2F\_5C A\_27a V0P)))$

Assume the following.

$$True \quad (3)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (4)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \quad (6)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (7)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}).((p (ap (c\_2Ebool\_2E\_3F\_21 A\_27a) (\lambda V1x \in A\_27a.(ap V0P V1x)))) \Leftrightarrow ((\exists V2x \in A\_27a.(p (ap V0P V2x)) \wedge (\forall V3x \in A\_27a.(\forall V4y \in A\_27a.((p (ap V0P V3x)) \wedge (p (ap V0P V4y)) \Rightarrow (V3x = V4y))))))) \quad (8)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow \forall A\_27c.nonempty A\_27c \Rightarrow (\forall V0f \in (A\_27b^{A\_27a}).(\forall V1g \in (A\_27c^{A\_27b}).(\forall V2x \in A\_27c.((ap (ap (ap (c\_2Ecombin\_2Eo A\_27c A\_27b A\_27a) V0f) V1g) V2x) = (ap V0f (ap V1g V2x)))))) \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow \forall A\_27c.nonempty A\_27c \Rightarrow (\forall V0f \in (A\_27c^{A\_27a}).(\forall V1g \in (A\_27c^{A\_27b}). \\ & (p (ap (c\_2Ebool\_2E\_3F\_21 (A\_27c^{(ty\_2Esum\_2Esum A\_27a A\_27b)})) \\ & (\lambda V2h \in (A\_27c^{(ty\_2Esum\_2Esum A\_27a A\_27b)}).(ap (ap c\_2Ebool\_2E\_2F\_5C \\ & (ap (ap (c\_2Emin\_2E\_3D (A\_27c^{A\_27a})) (ap (ap (c\_2Ecombin\_2Eo A\_27a \\ & A\_27c (ty\_2Esum\_2Esum A\_27a A\_27b)) V2h) (c\_2Esum\_2EINL A\_27a \\ & A\_27b))) V0f) (ap (ap (c\_2Emin\_2E\_3D (A\_27c^{A\_27b})) (ap (ap (c\_2Ecombin\_2Eo \\ & A\_27b A\_27c (ty\_2Esum\_2Esum A\_27a A\_27b)) V2h) (c\_2Esum\_2EINR \\ & A\_27a A\_27b))) V1g)))))) \quad (10) \end{aligned}$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow \forall A\_27c. \\ & nonempty\ A\_27c \Rightarrow (\forall V0f \in (A\_27c^{A\_27a}). (\forall V1g \in (A\_27c^{A\_27b}). \\ & (\exists V2h \in (A\_27c^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)}). ((\forall V3x \in \\ & A\_27a. ((ap\ V2h\ (ap\ (c\_2Esum\_2EINL\ A\_27a\ A\_27b)\ V3x)) = (ap\ V0f\ V3x)))) \wedge \\ & (\forall V4y \in A\_27b. ((ap\ V2h\ (ap\ (c\_2Esum\_2EINR\ A\_27a\ A\_27b)\ V4y)) = \\ & (ap\ V1g\ V4y)))))) \end{aligned}$$