

thm\_2Esum\_2Esum\_\_CASES  
(TMKHk6ZhM7zoHTVZ3TDpW9vSSh4MdsLWfmS)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 5** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2$

**Definition 6** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 7** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A$

**Definition 8** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 9** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

**Definition 10** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Esum\_2Esum A0 A1) \tag{1}$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Esum\_2EABS\_sum A\_27a A\_27b \in ((ty\_2Esum\_2Esum A\_27a A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \tag{2}$$

**Definition 11** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27b.(ap (c\_2Esum\_2EABS$

**Definition 12** We define  $c\_Esum\_2EINL$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27a.(ap (c\_Esum\_2EABS$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}). (\forall V1t \in \\ & A\_27a. ((\forall V2x \in A\_27a. ((V2x = V1t) \Rightarrow (p (ap\ V0P\ V2x)))) \Rightarrow (p ( \\ & ap (c\_Ebool\_2E\_3F\ A\_27a)\ V0P)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ & \forall V0P \in (2^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)}). (((\forall V1x \in \\ & A\_27a. (p (ap\ V0P (ap (c\_Esum\_2EINL\ A\_27a\ A\_27b)\ V1x)))) \wedge (\forall V2y \in \\ & A\_27b. (p (ap\ V0P (ap (c\_Esum\_2EINR\ A\_27a\ A\_27b)\ V2y)))))) \Rightarrow (\forall V3s \in \\ & (ty\_2Esum\_2Esum\ A\_27a\ A\_27b). (p (ap\ V0P\ V3s)))))) \end{aligned} \quad (4)$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ & \forall V0ss \in (ty\_2Esum\_2Esum\ A\_27a\ A\_27b). ((\exists V1x \in A\_27a. \\ & (V0ss = (ap (c\_Esum\_2EINL\ A\_27a\ A\_27b)\ V1x))) \vee (\exists V2y \in A\_27b. \\ & (V0ss = (ap (c\_Esum\_2EINR\ A\_27a\ A\_27b)\ V2y)))))) \end{aligned}$$