

thm_2Esum_2Esum__distinct
(TMdXQrNPR71mk9eaxEvfUmQcfrnGt8QkmYu)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Definition 5 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Esum_2Esum A0 A1) \tag{1}$$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Esum_2EABS_sum A_27a A_27b \in ((ty_2Esum_2Esum A_27a A_27b)^{((2^{A-27b})^{A-27a})^2}) \tag{2}$$

Definition 8 We define c_2Esum_2EINR to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27b.(ap (c_2Esum_2EABS_sum$

Definition 9 We define c_2Esum_2EINL to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27a.(ap (c_2Esum_2EABS_sum$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge p (ap P x)))$ of type $\iota \Rightarrow \iota$.

Definition 11 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40$

Assume the following.

$$True \quad (3)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (5)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\forall V0t1 \in A_27a. (\forall V1t2 \in A_27b. ((ap (\lambda V2x \in A_27b. V0t1) V1t2) = V0t1))) \quad (6)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \quad (7)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg (p V0t)) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg (p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg (p V0t)))))) \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow \forall A_27c. \\ & nonempty A_27c \Rightarrow (\forall V0f \in (A_27c^{A_27a}). (\forall V1g \in (A_27c^{A_27b}). \\ & (\exists V2h \in (A_27c^{ty_2Esum_2Esum A_27a A_27b}). ((\forall V3x \in \\ & A_27a. ((ap V2h (ap (c.2Esum_2EINL A_27a A_27b) V3x)) = (ap V0f V3x))) \wedge \\ & (\forall V4y \in A_27b. ((ap V2h (ap (c.2Esum_2EINR A_27a A_27b) V4y)) = \\ & (ap V1g V4y))))))) \end{aligned} \quad (10)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\ & \forall V0x \in A_27a. (\forall V1y \in A_27b. (\neg ((ap (c.2Esum_2EINL \\ & A_27a A_27b) V0x) = (ap (c.2Esum_2EINR A_27a A_27b) V1y)))) \end{aligned}$$