

thm\_2Etc\_2EDRESTR\_EMPTY  
(TMJ4vtWGbj7JT6EAKHGv4T5CdU9uVnBZiQe)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A-27a}). (ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})))$

**Definition 5** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

**Definition 6** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 7** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2. (ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

**Definition 8** We define  $c\_2Erelation\_2EEMPTY\_REL$  to be  $\lambda A\_27a : \iota. \lambda V0x \in A\_27a. \lambda V1y \in A\_27a. c\_2Ebool\_2E\_7E$

**Definition 9** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. c\_2Ebool\_2EF)$ .

**Definition 10** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. (\lambda V1f \in (2^{A-27a}). (ap V1f V0x)))$

**Definition 11** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap P x)) \mathbf{then} (the (\lambda x. x \in A \wedge p (ap P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 12** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A\_27a. (\lambda V2t2 \in A\_27a. (ap (c\_2Ebool\_2ECOND 2) (\lambda V3t3 \in 2.V3t3))$

**Definition 13** We define  $c\_2Etc\_2E\_5E\_7C$  to be  $\lambda A\_27a : \iota. \lambda V0R \in ((2^{A-27a})^{A-27a}). \lambda V1s \in (2^{A-27a}). \lambda V2a$

Assume the following.

$$True \quad (1)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (2)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (3)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t))))) \quad (5)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0t1 \in A\_27a. (\forall V1t2 \in A\_27a. ((ap (ap (ap (c\_2Ebool\_2ECOND A\_27a) c\_2Ebool\_2ET) V0t1) V1t2) = V0t1) \wedge ((ap (ap (ap (c\_2Ebool\_2ECOND A\_27a) c\_2Ebool\_2EF) V0t1) V1t2) = V1t2)))) \quad (6)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}). (\forall V1x \in A\_27a. ((p (ap (ap (c\_2Ebool\_2EIN A\_27a) V1x) V0P)) \Leftrightarrow (p (ap V0P V1x)))))) \quad (7)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a. (\neg(p (ap (ap (c\_2Ebool\_2EIN A\_27a) V0x) (c\_2Epred\_set\_2EEMPTY A\_27a)))))) \quad (8)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0R \in ((2^{A\_27a})^{A\_27a}). (\forall V1s \in (2^{A\_27a}). (\forall V2a \in A\_27a. ((ap (ap (ap (c\_2Etc\_2E\_5E\_7C A\_27a) V0R) V1s) V2a) = (ap (ap (ap (c\_2Ebool\_2ECOND (2^{A\_27a})) (ap (ap (c\_2Ebool\_2EIN A\_27a) V2a) V1s)) (ap V0R V2a)) (c\_2Epred\_set\_2EEMPTY A\_27a)))))) \quad (9)$$

**Theorem 1**

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0R \in ((2^{A\_27a})^{A\_27a}). ((ap (ap (c\_2Etc\_2E\_5E\_7C A\_27a) V0R) (c\_2Epred\_set\_2EEMPTY A\_27a)) = (c\_2Erelation\_2EEMPTY\_REL A\_27a)))$$