

# thm\_2Etc\_2EREMPTY\_\_RRESTR (TMVUcH9nYw3kSz6Fh32MdRdfMW31H2AynwF)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p \Rightarrow q)$  of type  $\iota$ .

**Definition 2** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Ebool_2E_2T` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

**Definition 4** We define `c_2Ebool_2E_21` to be  $\lambda A_{27a} : \iota. (\lambda V0P \in (2^{A_{27a}}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A_{27a}}))$

**Definition 5** We define `c_2Ebool_2E_2F` to be  $(\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V0t \in 2. V0t))$ .

**Definition 6** We define `c_2Ebool_2E_27E` to be  $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D\_3D\_3E } V0t) \text{c\_2Ebool\_2E\_2F}))$

**Definition 7** We define `c_2ERelation_2EEMPTY__REL` to be  $\lambda A_{27a} : \iota. \lambda V0x \in A_{27a}. \lambda V1y \in A_{27a}. \text{c\_2Ebool\_2E\_21}$

**Definition 8** We define `c_2Ebool_2E_2IN` to be  $\lambda A_{27a} : \iota. (\lambda V0x \in A_{27a}. (\lambda V1f \in (2^{A_{27a}}). (\text{ap } V1f V0x)))$

**Definition 9** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2) (\lambda V2t \in 2. V2t))$

**Definition 10** We define `c_2Etc_2E_27C_5E` to be  $\lambda A_{27a} : \iota. \lambda V0R \in ((2^{A_{27a}})^{A_{27a}}). \lambda V1s \in (2^{A_{27a}}). \lambda V2a$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p \Rightarrow V0t1) \Rightarrow (p \Rightarrow V1t2)) \Rightarrow (((p \Rightarrow V1t2) \Rightarrow (p \Rightarrow V0t1)) \Rightarrow ((p \Rightarrow V0t1) \Leftrightarrow (p \Rightarrow V1t2)))))) \tag{2}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((\text{True} \wedge (p \Rightarrow V0t)) \Leftrightarrow (p \Rightarrow V0t)) \wedge (((p \Rightarrow V0t) \wedge \text{True}) \Leftrightarrow \\ & (p \Rightarrow V0t)) \wedge (((\text{False} \wedge (p \Rightarrow V0t)) \Leftrightarrow \text{False}) \wedge (((p \Rightarrow V0t) \wedge \text{False}) \Leftrightarrow \text{False}) \wedge \\ & (((p \Rightarrow V0t) \wedge (p \Rightarrow V0t)) \Leftrightarrow (p \Rightarrow V0t)))))) \end{aligned} \tag{3}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t))))) \quad (5)$$

**Theorem 1**

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0s \in (2^{A_{.27a}}).((ap (ap (c_{.2Etc}_{.2E_{.7C}_{.5E}} A_{.27a}) (c_{.2Erelation}_{.2EEMPTY\_REL} A_{.27a})) V0s) = (c_{.2Erelation}_{.2EEMPTY\_REL} A_{.27a})))$$