

thm_2Etc_2ESUBSET__FDOM__LEM (TMQJ9JR1dqV4VbQDjsPMwjWjfbhBjTYzmgW)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow P \Rightarrow Q)$ of type ι .

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty \ ty_2Eone_2Eone \tag{1}$$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty \ A0 \Rightarrow \forall A1.nonempty \ A1 \Rightarrow nonempty \ (ty_2Esum_2Esum \ A0 \ A1) \tag{2}$$

Let $ty_2Efinite_map_2Efmap : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty \ A0 \Rightarrow \forall A1.nonempty \ A1 \Rightarrow nonempty \ (ty_2Efinite_map_2Efmap \ A0 \ A1) \tag{3}$$

Let $c_2Efinite_map_2Efmap_REP : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty \ A_27a \Rightarrow \forall A_27b.nonempty \ A_27b \Rightarrow c_2Efinite_map_2Efmap_REP \ A_27a \ A_27b \in (((ty_2Esum_2Esum \ A_27b \ ty_2Eone_2Eone)^{A_27a})(ty_2Efinite_map_2Efmap \ A_27a \ A_27b)) \tag{4}$$

Let $c_2Esum_2EISL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty \ A_27a \Rightarrow \forall A_27b.nonempty \ A_27b \Rightarrow c_2Esum_2EISL \ A_27a \ A_27b \in (2^{(ty_2Esum_2Esum \ A_27a \ A_27b)}) \tag{5}$$

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2ET to be $(ap \ (ap \ (c_2Emin_2E_3D \ (2^2)) \ (\lambda V0x \in 2.V0x)) \ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap \ (ap \ (c_2Emin_2E_3D \ (2^{A_27a})))$

Definition 5 We define $c_2Efinite_map_2EFDOM$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (ty_2Efinite_map_2Efmap \ A_27a \ A_27b)$

Definition 6 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 7 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2EF)$.

Let $c_2Esum_2EOUTL : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EOUTL \\ A_27a\ A_27b \in (A_27a^{(ty_2Esum_2Esum\ A_27a\ A_27b)}) \end{aligned} \quad (6)$$

Definition 8 We define $c_2Efinite_map_2EFAPPLY$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (ty_2Efinite_map_2EFmap\ A_27a\ (2^{A_27a}))$.

Definition 9 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap\ V1f\ V0x)))$.

Definition 10 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$.

Definition 11 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x.x \in A \wedge P\ x)) \text{ of type } \iota \Rightarrow \iota$.

Definition 12 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A_27a.(\lambda V2t2 \in A_27a.(\lambda V3t3 \in 2.V3t3))))$.

Definition 13 We define $c_2Etc_2EFMAP_TO_RELN$ to be $\lambda A_27a : \iota.\lambda V0f \in (ty_2Efinite_map_2EFmap\ A_27a\ (2^{A_27a}))$.

Definition 14 We define $c_2Epred_set_2ESUBSET$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).\lambda V1t \in (2^{A_27a}).(ap\ (c_2Epred_set_2EEMPTY\ A_27a)\ V1t))$.

Definition 15 We define $c_2Etc_2E_5E_7C$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).\lambda V1s \in (2^{A_27a}).\lambda V2a \in (2^{A_27a}).(\lambda V3x \in 2.V3x))$.

Definition 16 We define $c_2Etc_2E_7C_5E$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).\lambda V1s \in (2^{A_27a}).\lambda V2a \in (2^{A_27a}).(\lambda V3x \in 2.V3x))$.

Definition 17 We define $c_2Etc_2E_5E_7C_5E$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).\lambda V1s \in (2^{A_27a}).(\lambda V2a \in (2^{A_27a}).(\lambda V3x \in 2.V3x))$.

Definition 18 We define $c_2Erelation_2ERTC$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).\lambda V1a \in A_27a.\lambda V2b \in A_27a.(\lambda V3x \in 2.V3x))$.

Definition 19 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap\ V0P\ (ap\ (c_2Emin_2E_40\ A_27a)\ V0P))))$.

Definition 20 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.V2t))))$.

Definition 21 We define $c_2Etc_2EsubTC$ to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).\lambda V1s \in (2^{A_27a}).\lambda V2x \in (2^{A_27a}).(\lambda V3y \in 2.V3y))$.

Definition 22 We define $c_2Erelation_2ERDOM$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0R \in ((2^{A_27b})^{A_27a}).\lambda V1x \in A_27a.(\lambda V2y \in A_27b.(\lambda V3z \in 2.V3z))$.

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in \\ A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0f \in (ty_2Efinite_map_2EFmap \\ A_27a\ (2^{A_27a})).(p\ (ap\ (ap\ (c_2Epred_set_2ESUBSET\ A_27a)\ (ap \\ (c_2Erelation_2ERDOM\ A_27a\ A_27a)\ (ap\ (c_2Etc_2EFMAP_TO_RELN \\ A_27a)\ V0f))))\ (ap\ (c_2Efinite_map_2EFDOM\ A_27a\ (2^{A_27a})\ V0f)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0R \in ((2^{A_27a})^{A_27a}). \\
& (\forall V1s \in (2^{A_27a}). ((ap\ (c_2Erelation_2ERDOM\ A_27a\ A_27a) \\
& (ap\ (ap\ (c_2Etc_2EsubTC\ A_27a)\ V0R)\ V1s)) = (ap\ (c_2Erelation_2ERDOM \\
& \quad A_27a\ A_27a)\ V0R)))) \\
& \tag{9}
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0R \in ((2^{A_27a})^{A_27a}). \\
& (\forall V1s \in (2^{A_27a}). (\forall V2f \in (ty_2Efinite_map_2Efmap \\
& \quad A_27a\ (2^{A_27a})). (((ap\ (ap\ (c_2Etc_2EsubTC\ A_27a)\ V0R)\ V1s) = (\\
& \quad ap\ (c_2Etc_2EFMAP_TO_RELN\ A_27a)\ V2f)) \Rightarrow (p\ (ap\ (ap\ (c_2Epred_set_2ESUBSET \\
& \quad A_27a)\ (ap\ (c_2Erelation_2ERDOM\ A_27a\ A_27a)\ V0R))\ (ap\ (c_2Efinite_map_2EFDOM \\
& \quad \quad A_27a\ (2^{A_27a}))\ V2f))))))
\end{aligned}$$