

# thm\_2Etopology\_2Eclosed\_\_topspace (TMVAp- MDcewwB1mRPEdaxJbtkZ1NQB7F5rRM)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Ebool_2EIN` to be  $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(\lambda V1f V0x)))$

**Definition 4** We define `c_2Ebool_2EET` to be  $(\lambda p (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 5** We define `c_2Ebool_2E_21` to be  $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(\lambda p (ap (c_2Emin_2E_3D (2^{A_27a}))))$

**Definition 6** We define `c_2Ebool_2E_2F` to be  $(\lambda p (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 7** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

**Definition 8** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

Let `ty_2Epair_2Eprod` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty_2Epair_2Eprod A0 A1) \tag{1}$$

Let `c_2Epair_2EABS_prod` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS\_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \tag{2}$$

**Definition 9** We define `c_2Epair_2E_2C` to be  $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2Epair_2EABS\_prod A_27a A_27b) (ty_2Epair_2Eprod V0x V1y))$

Let `c_2Epred_set_2EGSPEC` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epred\_set\_2EGSPEC A_27a A_27b \in ((2^{A_27a})^{(ty_2Epair_2Eprod A_27a 2)^{A_27b}}) \tag{3}$$

**Definition 10** We define  $c\_2Epred\_set\_2EDIFF$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A-27a}).\lambda V1t \in (2^{A-27a}).(ap (c\_2Etopology\_2Etopology) A0)$ .  
Let  $ty\_2Etopology\_2Etopology : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Etopology\_2Etopology A0) \quad (4)$$

Let  $c\_2Etopology\_2Eopen\_in : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Etopology\_2Eopen\_in A\_27a \in ((2^{(2^{A-27a})}) (ty\_2Etopology\_2Etopology A\_27a)) \quad (5)$$

**Definition 11** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.$ **if**  $(\exists x \in A.p (ap P x))$  **then**  $(the (\lambda x.x \in A \wedge p (ap P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 12** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c\_2Emin\_2E\_40) A\_27a)))$ .

**Definition 13** We define  $c\_2Epred\_set\_2EBIGUNION$  to be  $\lambda A\_27a : \iota.\lambda V0P \in (2^{(2^{A-27a})}).(ap (c\_2Epred\_set\_2E) A\_27a)$ .

**Definition 14** We define  $c\_2Etopology\_2Etopspace$  to be  $\lambda A\_27a : \iota.\lambda V0top \in (ty\_2Etopology\_2Etopology A\_27a)$ .

**Definition 15** We define  $c\_2Epred\_set\_2ESUBSET$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A-27a}).\lambda V1t \in (2^{A-27a}).(ap (c\_2Epred\_set\_2E) A\_27a)$ .

**Definition 16** We define  $c\_2Epred\_set\_2EUNIV$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2E\_3F)$ .

**Definition 17** We define  $c\_2Etopology\_2Eclosed\_in$  to be  $\lambda A\_27a : \iota.\lambda V0top \in (ty\_2Etopology\_2Etopology A\_27a)$ .

**Definition 18** We define  $c\_2Etopology\_2Eclosed$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (ty\_2Etopology\_2Etopology A\_27a)$ .

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0s \in (2^{A-27a}).((p (ap (ap (c\_2Epred\_set\_2ESUBSET A\_27a) (c\_2Epred\_set\_2EUNIV A\_27a)) V0s)) \Leftrightarrow (V0s = (c\_2Epred\_set\_2EUNIV A\_27a)))) \quad (6)$$

**Theorem 1**

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0top \in (ty\_2Etopology\_2Etopology A\_27a).((p (ap (c\_2Etopology\_2Eclosed A\_27a) V0top)) \Rightarrow ((ap (c\_2Etopology\_2Etopspace A\_27a) V0top) = (c\_2Epred\_set\_2EUNIV A\_27a))))$$