

# thm\_2Etoto\_2EStrongLinearOrderExists (TMNcvK7fALn8FarcFarAZeYjBjpfHMgJ9kZ)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_40` to be  $\lambda A.\lambda P \in 2^A.$ **if**  $(\exists x \in A.p (ap P x))$  **then** (the  $(\lambda x.x \in A \wedge p x)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 2** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Ebool_2E_2T` to be  $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define `c_2Ebool_2E_21` to be  $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

**Definition 5** We define `c_2Ebool_2E_2F` to be  $(ap (c_2Ebool_2E_21 2)) (\lambda V0t \in 2.V0t)$ .

**Definition 6** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 7** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

**Definition 8** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.V2t$

**Definition 9** We define `c_2Ebool_2E_3F` to be  $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a$

**Definition 10** We define `c_2Erelation_2EWF` to be  $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_21$

**Definition 11** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.V2t$

**Definition 12** We define `c_2Erelation_2Etrichotomous` to be  $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_21$

**Definition 13** We define `c_2Erelation_2Etransitive` to be  $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_21$

**Definition 14** We define `c_2Erelation_2Eirreflexive` to be  $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_21$

**Definition 15** We define `c_2Erelation_2EStrongOrder` to be  $\lambda A_27g : \iota.\lambda V0Z \in ((2^{A_27g})^{A_27g}).(ap (ap c_2Ebool_2E_21$

**Definition 16** We define `c_2Erelation_2EStrongLinearOrder` to be  $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (ap c_2Ebool_2E_21$

Assume the following.

$$True \quad (1)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (2)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (3)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2. ((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (7)$$

Assume the following.

$$(\forall V0A \in 2. (((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (8)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. ((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (9)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. ((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))) \quad (10)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\exists V0R \in ((2^{A.27a})^{A.27a}). ((p (ap (c.2Relation_2EStrongLinearOrder A.27a) V0R)) \wedge (p (ap (c.2Relation_2EWF A.27a) V0R)))) \quad (11)$$

**Theorem 1**

$$\forall A.27a.nonempty A.27a \Rightarrow (\exists V0R \in ((2^{A.27a})^{A.27a}). (p (ap (c.2Relation_2EStrongLinearOrder A.27a) V0R)))$$