

thm_2Etoto_2EcharOrd__gt__lem
 (TMUTz8EHZzgy5EmnmVboZHFC5vJ3pJE2K98)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap (ap (c_2Emin_2E_3D (2^{A_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p \ P \Rightarrow p \ Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2. (ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2. (ap (c_2Ebool_2E_7E V2t) c_2Ebool_2EF))))))$

Let $ty_2Estring_2Echar : \iota$ be given. Assume the following.

$$nonempty \ ty_2Estring_2Echar \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty \ ty_2Enum_2Enum \quad (2)$$

Let $c_2Estring_2ECHR : \iota$ be given. Assume the following.

$$c_2Estring_2ECHR \in (ty_2Estring_2Echar^{ty_2Enum_2Enum}) \quad (3)$$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in omega \quad (4)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{omega}) \quad (5)$$

Definition 22 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum.$

Let $c_2EternaryComparisons_2ELESS : \iota$ be given. Assume the following.

$$c_2EternaryComparisons_2ELESS \in ty_2EternaryComparisons_2Eordering \quad (11)$$

Definition 23 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A_27a. (\lambda V2t2 \in A_27a. ($

Let $c_2EternaryComparisons_2EEQUAL : \iota$ be given. Assume the following.

$$c_2EternaryComparisons_2EEQUAL \in ty_2EternaryComparisons_2Eordering \quad (12)$$

Definition 24 We define $c_2Etoto_2ETO_of_LinearOrder$ to be $\lambda A_27a : \iota. \lambda V0r \in ((2^{A_27a})^{A_27a}). \lambda V1x \in$

Let $c_2Estring_2EORD : \iota$ be given. Assume the following.

$$c_2Estring_2EORD \in (ty_2Enum_2Enum^{ty_2Estring_2Echar}) \quad (13)$$

Definition 25 We define $c_2Etoto_2EnumOrd$ to be $(ap (c_2Etoto_2ETO_of_LinearOrder ty_2Enum_2Enum))$

Definition 26 We define $c_2Etoto_2EcharOrd$ to be $\lambda V0a \in ty_2Estring_2Echar. \lambda V1b \in ty_2Estring_2Echar.$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. \\ & \forall V2p \in ty_2Enum_2Enum. (((p (ap (ap c_2Eprim_rec_2E_3C \\ & V0m) V1n)) \wedge (p (ap (ap c_2Eprim_rec_2E_3C V1n) V2p))) \Rightarrow (p (ap (ap \\ & c_2Eprim_rec_2E_3C V0m) V2p)))))) \end{aligned} \quad (14)$$

Assume the following.

$$True \quad (15)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (16)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg (p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg (\\ & p V0t))))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & (\forall V0r \in ty_2Enum_2Enum. ((p (ap (ap c_2Eprim_rec_2E_3C \\ & V0r) (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 \\ & (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 \\ & (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 \\ & (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO))))))))))) \Rightarrow \\ & ((ap c_2Estring_2EORD (ap c_2Estring_2ECHR V0r)) = V0r))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned}
 & \forall A_27a. \text{nonempty } A_27a \Rightarrow (\forall V0r \in ((2^{A_27a})^{A_27a}). \\
 & ((p (ap (c_2Erelation_2EStrongLinearOrder A_27a) V0r)) \Rightarrow (\forall V1x \in \\
 & A_27a. (\forall V2y \in A_27a. ((ap (ap (ap (c_2Etoto_2ETO_of_LinearOrder \\
 & A_27a) V0r) V1x) V2y) = c_2EternaryComparisons_2EGREATER) \Leftrightarrow (p \\
 & (ap (ap V0r V2y) V1x))))))) \\
 \end{aligned} \tag{19}$$

Assume the following.

$$(p (ap (c_2Erelation_2EStrongLinearOrder ty_2Enum_2Enum) c_2Eprim_rec_2E_3C)) \tag{20}$$

Theorem 1

$$\begin{aligned}
 & (\forall V0a \in ty_2Enum_2Enum. (\forall V1b \in ty_2Enum_2Enum. (\\
 & ((ap (ap c_2Etoto_2EnumOrd V0a) V1b) = c_2EternaryComparisons_2EGREATER) \Rightarrow \\
 & ((p (ap (ap c_2Eprim_rec_2E_3C V0a) (ap c_2Earithmetic_2ENUMERAL \\
 & (ap c_2Earithmetic_2EBIT2 (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 \\
 & (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 \\
 & (ap c_2Earithmetic_2EBIT1 (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO))))))))))) \Leftrightarrow \\
 & \text{True}) \Rightarrow ((ap (ap c_2Etoto_2EcharOrd (ap c_2Estring_2ECHR V0a)) \\
 & (ap c_2Estring_2ECHR V1b)) = c_2EternaryComparisons_2EGREATER)))))) \\
 \end{aligned}$$