

# thm\_2Etoto\_2EcharOrd\_\_thm (TML- HiHQ8VUbAge8L5BMvbkLH6YgQpsdrJqH)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $ty\_2Estring\_2Echar : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Estring\_2Echar \tag{2}$$

Let  $c\_2Estring\_2EORD : \iota$  be given. Assume the following.

$$c\_2Estring\_2EORD \in (ty\_2Enum\_2Enum^{ty\_2Estring\_2Echar}) \tag{3}$$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda \tau a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap (c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{6}$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\mathbf{if}\ (\exists x \in A.p\ (ap\ P\ x))\ \mathbf{then}\ (the\ (\lambda x.x \in A \wedge p$   
of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

**Definition 12** We define  $c\_2Estring\_2Echar\_lt$  to be  $\lambda V0a \in ty\_2Estring\_2Echar.\lambda V1b \in ty\_2Estring\_2E$

Let  $ty\_2EternaryComparisons\_2Eordering : \iota$  be given. Assume the following.

$$nonempty\ ty\_2EternaryComparisons\_2Eordering \quad (7)$$

Let  $c\_2EternaryComparisons\_2EGREATER : \iota$  be given. Assume the following.

$$c\_2EternaryComparisons\_2EGREATER \in ty\_2EternaryComparisons\_2Eordering \quad (8)$$

Let  $c\_2EternaryComparisons\_2ELESS : \iota$  be given. Assume the following.

$$c\_2EternaryComparisons\_2ELESS \in ty\_2EternaryComparisons\_2Eordering \quad (9)$$

Let  $c\_2EternaryComparisons\_2EEQUAL : \iota$  be given. Assume the following.

$$c\_2EternaryComparisons\_2EEQUAL \in ty\_2EternaryComparisons\_2Eordering \quad (10)$$

**Definition 13** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A\_27a.(\lambda V2t2 \in A\_27a.($

**Definition 14** We define  $c\_2Etoto\_2ETO\_of\_LinearOrder$  to be  $\lambda A\_27a : \iota.\lambda V0r \in ((2^{A\_27a})^{A\_27a}).\lambda V1x \in$

**Definition 15** We define  $c\_2Etoto\_2EnumOrd$  to be  $(ap\ (c\_2Etoto\_2ETO\_of\_LinearOrder\ ty\_2Enum\_2Enum$

**Definition 16** We define  $c\_2Etoto\_2EcharOrd$  to be  $\lambda V0a \in ty\_2Estring\_2Echar.\lambda V1b \in ty\_2Estring\_2E$

Assume the following.

$$True \quad (11)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (12)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (13)$$

Assume the following.

$$(\forall V0a \in ty\_2Estring\_2Echar.(\forall V1a\_27 \in ty\_2Estring\_2Echar.(((ap\ c\_2Estring\_2EORD\ V0a) = (ap\ c\_2Estring\_2EORD\ V1a\_27)) \Leftrightarrow (V0a = V1a\_27)))) \quad (14)$$

**Theorem 1**

$$(c\_2Etoto\_2EcharOrd = (ap (c\_2Etoto\_2ETO\_of\_LinearOrder ty\_2Estring\_2Echar) c\_2Estring\_2Echar\_lt))$$