

thm_2Etoto_2Etrichotomous__ALT (TMbxxAQJVbyxp8cv7RJ1cZLqbFV3g4VrYBc)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 7 We define `c_2Ebool_2E_27E` to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Definition 8 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Definition 9 We define `c_2Erelation_2Etrichotomous` to be $\lambda A_27a : \iota.\lambda V0R \in ((2^{A_27a})^{A_27a}).(ap (c_2Ebool_2E_27E$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t) \Leftrightarrow (p V1x))) \tag{2}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \tag{4}$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (5)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (\forall V2C \in 2. (((p\ V0A) \vee (p\ V1B) \vee (p\ V2C)) \Leftrightarrow (((p\ V0A) \vee (p\ V1B)) \vee (p\ V2C)))))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p\ V0A) \wedge (p\ V1B))) \Leftrightarrow ((\neg(p\ V0A) \vee \neg(p\ V1B))) \wedge ((\neg((p\ V0A) \vee (p\ V1B))) \Leftrightarrow ((\neg(p\ V0A) \wedge \neg(p\ V1B)))))))) \quad (7)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((p\ V0A) \Rightarrow (p\ V1B)) \Leftrightarrow ((\neg(p\ V0A) \vee (p\ V1B)))))) \quad (8)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0R \in ((2^{A_27a})^{A_27a}). ((p\ (ap\ (c_2Erelation_2Etrichotomous\ A_27a)\ V0R)) \Leftrightarrow (\forall V1x \in A_27a. (\forall V2y \in A_27a. (((\neg(p\ (ap\ (ap\ V0R\ V1x)\ V2y))) \wedge (\neg(p\ (ap\ (ap\ V0R\ V2y)\ V1x)))) \Rightarrow (V1x = V2y))))))$$