

thm_2Etrasc_2ESQRT__0
(TMXv4w21Dor9ditVoWKx8pyir5eezmWYDEZ)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))\ a))$

Definition 5 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (c_2Enum_2ESUC_REP\ m))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 6 We define $c_Earithmic_EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_Earithmic_$

Definition 7 We define $c_Earithmic_EZERO$ to be c_2Enum_2E0 .

Definition 8 We define $c_Earithmic_EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_Earithmic_$

Definition 9 We define $c_Earithmic_ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \quad (7)$$

Let $c_2Ereal_2Epow : \iota$ be given. Assume the following.

$$c_2Ereal_2Epow \in ((ty_2Erealax_2Ereal^{ty_2Enum_2Enum})^{ty_2Erealax_2Ereal}) \quad (8)$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (9)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \quad (10)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (11)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax_2Ereal}) \quad (12)$$

Definition 10 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\mathbf{if} (\exists x \in A.p (ap\ P\ x)) \mathbf{then} (the (\lambda x.x \in A \wedge p\ x))$ of type $\iota \Rightarrow \iota$.

Definition 11 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap (c_2Emin_2E_40 (t$

Let $c_2Erealax_2Etreallt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etreallt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \quad (13)$$

Definition 12 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 13 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 14 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21\ 2) (\lambda V2t \in 2$

Definition 15 We define $c_Etransc_Eroot$ to be $\lambda V0n \in ty_Eenum_Eenum. \lambda V1x \in ty_Erealx_Ereal$

Definition 16 We define $c_Etransc_Esqrt$ to be $\lambda V0x \in ty_Erealx_Ereal. (ap (ap c_Etransc_Eroot ($

Assume the following.

$$\begin{aligned} & ((ap c_Earithmic_EENUMERAL (ap c_Earithmic_EEBIT2 c_Earithmic_EZERO)) = \\ & (ap c_Eenum_ESUC (ap c_Earithmic_EENUMERAL (ap c_Earithmic_EEBIT1 \\ & c_Earithmic_EZERO)))) \end{aligned} \tag{14}$$

Assume the following.

$$True \tag{15}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \tag{16}$$

Assume the following.

$$\begin{aligned} & (\forall V0n \in ty_Eenum_Eenum. ((ap (ap c_Etransc_Eroot (ap c_Eenum_ESUC \\ & V0n)) (ap c_Ereal_Ereal_of_num c_Eenum_E0)) = (ap c_Ereal_Ereal_of_num \\ & c_Eenum_E0))) \end{aligned} \tag{17}$$

Theorem 1

$$\begin{aligned} & ((ap c_Etransc_Esqrt (ap c_Ereal_Ereal_of_num c_Eenum_E0)) = \\ & (ap c_Ereal_Ereal_of_num c_Eenum_E0)) \end{aligned}$$