

thm_2Etrasc_2ESQRT__MUL
(TMY3Uyz4cvq2WdWyqzjDSGXZaxLidTtQfpg)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V1x \in 2.V1x)))$

Definition 5 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (c_2Enum_2E0\ m))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 6 We define $c_Earithmic_EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_Earithmic_E$

Definition 7 We define c_Ebool_EF to be $(ap (c_Ebool_E21 2) (\lambda V0t \in 2.V0t))$.

Definition 8 We define $c_Emin_E3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 9 We define c_Ebool_E7E to be $(\lambda V0t \in 2.(ap (ap c_Emin_E3D_3D_3E V0t) c_Ebool_EF$

Definition 10 We define $c_Earithmic_EZERO$ to be c_2Enum_2E0 .

Definition 11 We define $c_Earithmic_EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_Earithmic_E$

Definition 12 We define $c_Earithmic_ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{7}$$

Let $c_2Ereal_2Epow : \iota$ be given. Assume the following.

$$c_2Ereal_2Epow \in ((ty_2Erealx_2Ereal^{ty_2Enum_2Enum})^{ty_2Erealx_2Ereal}) \tag{8}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{9}$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{10}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{11}$$

Let $c_2Erealx_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealx_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealx_2Ereal}) \tag{12}$$

Definition 13 We define c_Emin_E40 to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge P x)$ of type $\iota \Rightarrow \iota$).

Definition 14 We define $c_2Erealx_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealx_2Ereal.(ap (c_Emin_E40 (t$

Let $c_2Erealx_2Etreal_lt : \iota$ be given. Assume the following.

$$c_2Erealx_2Etreal_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)}) \tag{13}$$

Definition 15 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 16 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21) 2) (\lambda V2t \in 2.))$

Definition 17 We define $c_2Etransc_2Eroot$ to be $\lambda V0n \in ty_2Enum_2Enum.\lambda V1x \in ty_2Erealax_2Ereal.$

Definition 18 We define $c_2Etransc_2Esqrt$ to be $\lambda V0x \in ty_2Erealax_2Ereal.(ap (ap c_2Etransc_2Eroot ($

Let $c_2Erealax_2Etrealmul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmul \in (((ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)))(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal) \quad (14)$$

Let $c_2Erealax_2Etrealeq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealeq \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)) \quad (15)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal)^{(2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)})(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)} \quad (16)$$

Definition 19 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)$

Definition 20 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal.$

Definition 21 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal.$

Assume the following.

$$((ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO)) = (ap c_2Enum_2ESUC (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO)))) \quad (17)$$

Assume the following.

$$True \quad (18)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A.27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (19)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(p V0t)))))) \quad (20)$$

Assume the following.

$$\begin{aligned}
& (\forall V0n \in ty_2Enum_2Enum. (\forall V1x \in ty_2Erealx_2Ereal. \\
& (\forall V2y \in ty_2Erealx_2Ereal. (((p (ap (ap c_2Ereal_2Ereal_lte \\
& (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V1x)) \wedge (p (ap (ap c_2Ereal_2Ereal_lte \\
& (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V2y)))) \Rightarrow ((ap (ap c_2Etrasc_2Eroot \\
& (ap c_2Enum_2ESUC V0n)) (ap (ap c_2Erealx_2Ereal_mul V1x) V2y)) = \\
& (ap (ap c_2Erealx_2Ereal_mul (ap (ap c_2Etrasc_2Eroot (ap c_2Enum_2ESUC \\
& V0n)) V1x)) (ap (ap c_2Etrasc_2Eroot (ap c_2Enum_2ESUC V0n)) V2y))))))))) \\
& \hspace{15em} (21)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& (\forall V0x \in ty_2Erealx_2Ereal. (\forall V1y \in ty_2Erealx_2Ereal. \\
& (((p (ap (ap c_2Ereal_2Ereal_lte (ap c_2Ereal_2Ereal_of_num \\
& c_2Enum_2E0)) V0x)) \wedge (p (ap (ap c_2Ereal_2Ereal_lte (ap c_2Ereal_2Ereal_of_num \\
& c_2Enum_2E0)) V1y)))) \Rightarrow ((ap c_2Etrasc_2Esqrt (ap (ap c_2Erealx_2Ereal_mul \\
& V0x) V1y)) = (ap (ap c_2Erealx_2Ereal_mul (ap c_2Etrasc_2Esqrt \\
& V0x)) (ap c_2Etrasc_2Esqrt V1y)))))))))
\end{aligned}$$