

thm_2Etransc_2ESQRT__POW_2
 (TMTxYaoDMm59c96VD8WfqHM1P7AmqfkyAkx)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A. \lambda P \in (2^{A_27a}). (ap (ap (c_2Emin_2E_3D (2^{A_27a})) (\lambda V0P \in (2^{A_27a})) (\lambda V1P \in (2^{A_27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2. (ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2. (ap (c_2Ebool_2E_7E V2t) c_2Ebool_2EF))))))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in omega \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (2)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{omega}) \quad (3)$$

Definition 8 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \quad (4)$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealax_2Ereal^{ty_2Enum_2Enum}) \quad (5)$$

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

nonempty $ty_2Ehreal_2Ehreal$ (6)

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.\text{nonempty } A0 \Rightarrow \forall A1.\text{nonempty } A1 \Rightarrow \text{nonempty}(\text{ty_2Epair_2Eprod } A0\ A1)$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{ty_2Erealax}) \\ (8)$$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p (ap P x))$ then $(the (\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_2\text{Erealax_2Ereal_REP}$ to be $\lambda V0a \in ty_2\text{Erealax_2Ereal}.(ap(c_2\text{Emin_2E_40}(t\backslash a)))$

Let $c_2Erealax_2Etreal_lt : \iota$ be given. Assume the following.

$$c_2 Erealax_2Etreal_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)})^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)} \quad (9)$$

Definition 11 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal. \lambda V1T2 \in ty_2Erealax_2Ereal. \dots$

Definition 12 We define $c_2Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Er$

Definition 13 We define $c_2Earthmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (10)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^\omega)^\omega \quad (11)$$

Definition 14 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \\ (12)$$

Definition 15 We define $c_2Earthmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap (ap c_2Earthmetic$

Definition 16 We define `c_2Earithmetic_2ENUMERAL` to be $\lambda V0x \in ty_Enum_2Enum. V0x$.

Let $c_2Ereal_2Epow : \iota$ be given. Assume the following.

$$c_2Ereal_2Epow \in ((ty_2Erealax_2Ereal^{ty_2Enum_2Enum})^{ty_2Erealax_2Ereal}) \quad (13)$$

Definition 17 We define $c_2Etransc_2Eroot$ to be $\lambda V0n \in ty_2Enum_2Enum. \lambda V1x \in ty_2Erealax_2Ereal.$

Definition 18 We define $c_2Etransc_2Esqrt$ to be $\lambda V0x \in ty_2Erealax_2Ereal. (ap (ap c_2Etransc_2Eroot ($

Assume the following.

$$True \quad (14)$$

Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (15)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t))))))) \quad (16)$$

Assume the following.

$$(\forall V0x \in ty_2Erealax_2Ereal. (((ap (ap c_2Ereal_2Epow (ap c_2Etransc_2Esqrt V0x)) (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO))) = V0x) \Leftrightarrow (p (ap (ap c_2Ereal_2Ereal_lte (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V0x)))) \quad (17)$$

Theorem 1

$$(\forall V0x \in ty_2Erealax_2Ereal. ((p (ap (ap c_2Ereal_2Ereal_lte (ap c_2Ereal_2Ereal_of_num c_2Enum_2E0)) V0x)) \Rightarrow ((ap (ap c_2Ereal_2Epow (ap c_2Etransc_2Esqrt V0x)) (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT2 c_2Earithmetic_2EZERO))) = V0x)))$$