

thm_2Ewhile_2ELEAST_-T (TMXCd-
NAWakSZ6LZy7x6rnZV4RvZqMNWxdxt)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (2)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (3)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$.

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ (ap\ (c_2Emin_2E_3D\ (2^{A_27a}))\ (\lambda V1t \in 2.V1t)))$.

Definition 5 We define c_2Ebool_2EF to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (5)$$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ c_2Enum_2EABS_num\ m)$.

Definition 7 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2. \lambda Q \in 2. inj_o (p \Rightarrow p \ Q)$ of type ι .

Definition 8 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF))$

Definition 9 We define $c_2Ecombin_2Eo$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda A_27c : \iota.\lambda V0f \in (A_27b^{A_27c}).\lambda V1g$

Let $c_2Ewhile_2EWHILE : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Ewhile_2EWHILE A_27a \in (((A_27a^{A_27a})^{(A_27a^{A_27a})})^{(2^{A_27a})}) \quad (6)$$

Definition 10 We define $c_2Ewhile_2ELEAST$ to be $\lambda V0P \in (2^{ty_2Enum_2Enum}).(ap (ap (ap (c_2Ewhile_2EWHILE V0P)))$

Definition 11 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Definition 12 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (\text{the } (\lambda x.x \in A \wedge p \text{ of type } \iota \Rightarrow \iota).$

Definition 13 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 V0P)))$

Definition 14 We define $c_2Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2Enum.$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum.((\neg(V0n = c_2Enum_2E0)) \Leftrightarrow (p (ap (ap c_2Eprim_rec_2E_3C c_2Enum_2E0) V0n)))) \quad (7)$$

Assume the following.

$$True \quad (8)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\exists V1x \in A_27a.(p V0t)) \Leftrightarrow (p V0t))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (11)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (12)$$

Assume the following.

$$\forall A. \exists a. \text{nonempty } A \Rightarrow (\forall V0x \in A. \exists a. (\forall V1y \in A. \exists a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$(\forall V0t \in 2. (((\text{True} \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow \text{True}) \Leftrightarrow (p V0t)) \wedge (((\text{False} \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow \text{False}) \Leftrightarrow (\neg(p V0t))))))) \quad (14)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (15)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1x_27 \in 2. (\forall V2y \in 2. (\forall V3y_27 \in 2. (((p V0x) \Leftrightarrow (p V1x_27)) \wedge ((p V1x_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_27)))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_27) \Rightarrow (p V3y_27))))))) \quad (16)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (17)$$

Assume the following.

$$(\forall V0A \in 2. ((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow \text{False}))) \quad (18)$$

Assume the following.

$$(\forall V0A \in 2. (((\neg(p V0A)) \Rightarrow \text{False}) \Rightarrow (((p V0A) \Rightarrow \text{False}) \Rightarrow \text{False}))) \quad (19)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. ((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (20)$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. ((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))))) \quad (21)$$

Assume the following.

$$\begin{aligned} & (\forall V0Q \in (2^{ty_2Enum_2Enum}). (\forall V1P \in (2^{ty_2Enum_2Enum}). \\ & (((\exists V2n \in ty_2Enum_2Enum. (p (ap V1P V2n))) \wedge (\forall V3n \in \\ & ty_2Enum_2Enum. (((\forall V4m \in ty_2Enum_2Enum. ((p (ap (ap c_2Eprim_rec_2E_3C \\ & V4m) V3n)) \Rightarrow (\neg(p (ap V1P V4m)))) \wedge (p (ap V1P V3n)) \Rightarrow (p (ap V0Q V3n)))) \Rightarrow \\ & (p (ap V0Q (ap c_2Ewhile_2ELEAST V1P))))))) \end{aligned} \quad (22)$$

Theorem 1

$$((ap c_2Ewhile_2ELEAST (\lambda V0x \in ty_2Enum_2Enum. c_2Ebool_2ET)) = c_2Enum_2E0)$$