

thm_2Ewords_2EBITS_ZEROL_DIMINDEX
(TMNFrpryfkReYpJt3LYLFeSx9FqTEW577D5)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_7E$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (ap c_2Enum_2EREP_num (ap c_2Enum_2ESUC_REP m)))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define $c_Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap\ V0P\ (ap\ (c_Emin_2E_40$

Definition 11 We define $c_Eprim_rec_2E_3C$ to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{5}$$

Definition 12 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 13 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 14 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum. (ap\ (ap\ c_2Earithmetic$

Definition 15 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum. V0x$.

Let $ty_2Ebool_2Eitself : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. nonempty\ A0 \Rightarrow nonempty\ (ty_2Ebool_2Eitself\ A0) \tag{7}$$

Let $c_2EfcP_2Edimindex : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty\ A_27a \Rightarrow c_2EfcP_2Edimindex\ A_27a \in (ty_2Enum_2Enum^{(ty_2Ebool_2Eitself\ A_27a)}) \tag{8}$$

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{9}$$

Definition 16 We define $c_2Earithmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum. (ap\ (ap\ c_2Earithmetic$

Let $c_2Earithmetic_2EEXP : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EEXP \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{10}$$

Let $c_2Earithmetic_2EDIV : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EDIV \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{11}$$

Definition 17 We define $c_2Ebit_2EDIV_2EXP$ to be $\lambda V0x \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Let $c_2Earithmetic_2EMOD : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EMOD \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{12}$$

Definition 18 We define $c_2Ebit_2EMOD_2EXP$ to be $\lambda V0x \in ty_2Enum_2Enum.\lambda V1n \in ty_2Enum_2E$

Definition 19 We define c_2Ebit_2EBITS to be $\lambda V0h \in ty_2Enum_2Enum.\lambda V1l \in ty_2Enum_2Enum.\lambda V$

Let $c_2Ebool_2Ethe_value : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Ebool_2Ethe_value\ A_27a \in (ty_2Ebool_2Eitself\ A_27a) \quad (13)$$

Let $c_2Ewords_2Edimword : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Ewords_2Edimword\ A_27a \in (ty_2Enum_2Enum^{(ty_2Ebool_2Eitself\ A_27a)}) \quad (14)$$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum.(\forall V1k \in ty_2Enum_2Enum.(p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ V1k)\ V0n) \Rightarrow ((ap\ (ap\ c_2Earithmetic_2EMOD\ V1k)\ V0n) = V1k)))) \quad (15)$$

Assume the following.

$$True \quad (16)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t) \Leftrightarrow (p\ V0t))) \quad (17)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \quad (18)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (19)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (20)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg (p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \quad (21)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p\ V0t1) \Rightarrow ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (22)$$

Assume the following.

$$2.(((p \ V0x) \Leftrightarrow (p \ V1x_27)) \wedge ((p \ V1x_27) \Rightarrow ((p \ V2y) \Leftrightarrow (p \ V3y_27)))) \Rightarrow \quad (23)$$

$$(((p \ V0x) \Rightarrow (p \ V2y)) \Leftrightarrow ((p \ V1x_27) \Rightarrow (p \ V3y_27))))$$

Assume the following.

$$\forall A_27a.nonempty \ A_27a \Rightarrow (\forall V0n \in ty_2Enum_2Enum. ($$

$$(ap \ (ap \ c_2Earithmetic_2EMOD \ V0n) \ (ap \ (c_2Ewords_2Edimword \ A_27a)$$

$$(c_2Ebool_2Ethe_value \ A_27a))) = (ap \ (ap \ (ap \ c_2Ebit_2EBITS \ ($$

$$ap \ (ap \ c_2Earithmetic_2E_2D \ (ap \ (c_2Efcf_2Edimindex \ A_27a) \ (c_2Ebool_2Ethe_value$$

$$A_27a))) \ (ap \ c_2Earithmetic_2ENUMERAL \ (ap \ c_2Earithmetic_2EBIT1$$

$$c_2Earithmetic_2EZERO)))) \ c_2Enum_2E0) \ V0n))) \quad (24)$$

Theorem 1

$$\forall A_27a.nonempty \ A_27a \Rightarrow (\forall V0n \in ty_2Enum_2Enum. ($$

$$(p \ (ap \ (ap \ c_2Eprim_rec_2E_3C \ V0n) \ (ap \ (c_2Ewords_2Edimword \ A_27a)$$

$$(c_2Ebool_2Ethe_value \ A_27a)))) \Rightarrow ((ap \ (ap \ (ap \ c_2Ebit_2EBITS$$

$$(ap \ (ap \ c_2Earithmetic_2E_2D \ (ap \ (c_2Efcf_2Edimindex \ A_27a) \ ($$

$$c_2Ebool_2Ethe_value \ A_27a))) \ (ap \ c_2Earithmetic_2ENUMERAL$$

$$(ap \ c_2Earithmetic_2EBIT1 \ c_2Earithmetic_2EZERO)))) \ c_2Enum_2E0)$$

$$V0n) = V0n)))$$