

thm_2Ewords_2Edimindex__1
(TMM4uh8m2Ya2Yxr3zEVkThDjiysMnNkMKrh)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define `c_2Earithmetic_2EBIT1` to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmetic_2EBIT1$

Definition 8 We define `c_2Earithmetic_2ENUMERAL` to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let `ty_2Ebool_2Eitself` : $\iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Ebool_2Eitself A0) \quad (7)$$

Let `c_2Ebool_2Ethe_value` : $\iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Ebool_2Ethe_value A_27a \in (ty_2Ebool_2Eitself A_27a) \quad (8)$$

Let `ty_2Eone_2Eone` : ι be given. Assume the following.

$$nonempty ty_2Eone_2Eone \quad (9)$$

Let `c_2Efcf_2Edimindex` : $\iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Efcf_2Edimindex A_27a \in (ty_2Enum_2Enum^{(ty_2Ebool_2Eitself A_27a)}) \quad (10)$$

Assume the following.

$$((ap (c_2Efcf_2Edimindex ty_2Eone_2Eone) (c_2Ebool_2Ethe_value ty_2Eone_2Eone)) = (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO))) \quad (11)$$

Theorem 1

$$((ap (c_2Efcf_2Edimindex ty_2Eone_2Eone) (c_2Ebool_2Ethe_value ty_2Eone_2Eone)) = (ap c_2Earithmetic_2ENUMERAL (ap c_2Earithmetic_2EBIT1 c_2Earithmetic_2EZERO)))$$