

l10\_amistd\_1 (TMTbmN-  
ThXrC2ENGaVvzgxGaZKqFKcY5f1Re)

October 27, 2020

Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_setfam\_1 : \iota \Rightarrow o$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v2\_memstr\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v3\_memstr\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $l1\_extpro\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_amistd\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow ((X1 = k9\_finseq\_1 X0) \Leftrightarrow ((k3\_finseq\_1 X1 = np\_1) \wedge (k10\_xtuple\_0 X1 = k1\_tarski X0))) \quad (1)$$

Assume the following.

$$\forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 X0) \Rightarrow (k7\_partfun1 X0 (k12\_finseq\_1 X0 X1) np\_1 = X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2. (m2\_subset\_1 X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \quad (3)$$

Assume the following.

$$\forall X0. k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow (k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (6)$$

Assume the following.

$$\forall X0.v1\_finseq\_1 (k5\_finseq\_1 X0) \quad (7)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (8)$$

Assume the following.

$$\forall X0.\neg v1\_xboole\_0 (k5\_finseq\_1 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k9\_finseq\_1 X0))\wedge(v1\_funct\_1 (k9\_finseq\_1 X0)) \quad (10)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow (m2\_finseq\_1 (k12\_finseq\_1 X0 X1) X0) \quad (12)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))\Rightarrow(v1\_xboole\_0 X1)) \quad (13)$$

**Theorem 1**

$$\begin{aligned} &\forall X0.(m2\_subset\_1 X0 k1\_numbers k5\_numbers)\Rightarrow(\forall X1. \\ &(\neg v1\_setfam\_1 X1)\Rightarrow(\forall X2.((\neg v2\_struct\_0 X2)\wedge((v2\_memstr\_0 \\ &X2 X1)\wedge((v3\_memstr\_0 X2 X1)\wedge(l1\_extpro\_1 X2 X1))))\Rightarrow(\exists X3. \\ &((\neg v1\_xboole\_0 X3)\wedge(m2\_finseq\_1 X3 k5\_numbers))\wedge((k7\_partfun1 \\ &k5\_numbers X3 np\_1 = X0)\wedge((k7\_partfun1 k5\_numbers X3 (k3\_finseq\_1 \\ &X3) = X0)\wedge(\forall X4.(m2\_subset\_1 X4 k1\_numbers k5\_numbers)\Rightarrow \\ &((r1\_xxreal\_0 np\_1 X4)\Rightarrow((r1\_xxreal\_0 (k3\_finseq\_1 X3) X4)\vee( \\ &k7\_partfun1 k5\_numbers X3 (k2\_nat\_1 X4 np\_1) \in k3\_amistd\_1 X1 X2 \\ &(k7\_partfun1 k5\_numbers X3 X4)))))))))) \end{aligned}$$