

l10_fib_num

(TMJbeeMtT1SmgbxjNYHMddcMx5oWb1mTjcc)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_pre_ff : \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $np_0 : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg(k6_numbers \neq X0) \wedge (r1_xxreal_0 X0 k6_numbers)) \quad (2)$$

Assume the following.

$$\begin{aligned} & (k1_pre_ff k6_numbers = k6_numbers) \wedge ((k1_pre_ff np_1 = np_1) \wedge \\ & (\forall X0.(v7_ordinal1 X0) \Rightarrow (k1_pre_ff (k2_nat_1 (k1_nat_1 \\ & X0 np_1) np_1) = k2_nat_1 (k1_pre_ff X0) (k1_pre_ff (k1_nat_1 \\ & X0 np_1)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (5)$$

Assume the following.

$$k2_xcmplx_0 np_1 np_0 = np_1 \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(m1_subset_1 X1 k5_numbers))\Rightarrow(k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (10)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(r1_xxreal_0 X0 (k1_pre_ff (k1_nat_1 X0 np_1))) \quad (11)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_nat_1 X1 X0) \quad (13)$$

Assume the following.

$$\forall X0.(v6_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v7_ordinal1 X1)) \quad (14)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k5_numbers)\Rightarrow(k1_pre_ff (k2_nat_1 X0 np_1)\neq k6_numbers)$$