

l10\_gobrd11 (TMNvYC-  
qUggy3XmR1VA9PoyY8QYHtE7DiFzE)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k19\_euclid : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k1\_euclid : \iota \Rightarrow \iota$  be given. Let  $v6\_membered : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (2)$$

Assume the following.

$$((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$u1\_struct\_0 (k15\_euclid np\_2) = k1\_euclid np\_2 \quad (5)$$

Assume the following.

$$v6\_membered k4\_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0. (v7\_ordinal1 X0) \Rightarrow (\neg v1\_xboole\_0 (k1\_euclid X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(m1\_subset\_1 (k19\_euclid X0 X1) (u1\_struct\_0 (k15\_euclid np\_2))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(r1\_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (9)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v6\_membered X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 X0)\Rightarrow (v7\_ordinal1 X1)) \quad (11)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(m1\_subset\_1 (ReplSep2 (toset (\lambda X1 : \iota.m1\_subset\_1 X1 k1\_numbers)) (\lambda X1 : \iota.toset (\lambda X2 : \iota.m1\_subset\_1 X2 k1\_numbers)) (\lambda X1 : \iota.\lambda X2 : \iota.r1\_xreal\_0 X0 X2) (\lambda X1 : \iota.\lambda X2 : \iota.k19\_euclid X1 X2)) (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid np\_2))))$$