

# l114\_jordan (TMGWa- jCdF3p3hVuy5pnqRucn5zMDTt2RtV5)

October 27, 2020

Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_rltopsp1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k19\_euclid : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_real\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $np\_3 : \iota$  be given. Let  $k4\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $v6\_membered : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $l1\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. r1\_tarski X0 (k2\_xboole\_0 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_3) \wedge (m2\_subset\_1 np\_3 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_3 k5\_numbers) \wedge (m1\_subset\_1 np\_3 k1\_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)))\Rightarrow(k4\_subset\_1 X0 X1 X2 = k2\_xboole\_0 X1 X2) \quad (6)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(k1\_real\_1 X0 = k4\_xcmplx\_0 X0) \quad (7)$$

Assume the following.

$$v6\_membered k4\_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow(((v2\_pre\_topc (k15\_euclid X0))\wedge((v13\_algstr\_0 (k15\_euclid X0))\wedge((v2\_rlvect\_1 (k15\_euclid X0))\wedge((v3\_rlvect\_1 (k15\_euclid X0))\wedge((v4\_rlvect\_1 (k15\_euclid X0))\wedge((v5\_rlvect\_1 (k15\_euclid X0))\wedge((v6\_rlvect\_1 (k15\_euclid X0))\wedge((v7\_rlvect\_1 (k15\_euclid X0))\wedge((v8\_rlvect\_1 (k15\_euclid X0))\wedge(v5\_rltopsp1 (k15\_euclid X0)))))))))))))) \quad (9)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow((\neg v2\_struct\_0 (k15\_euclid X0))\wedge(v5\_rltopsp1 (k15\_euclid X0))) \quad (10)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow((v1\_xcmplx\_0 (k4\_xcmplx\_0 X0))\wedge(v1\_xreal\_0 (k4\_xcmplx\_0 X0))) \quad (11)$$

Assume the following.

$$\forall X0.(l1\_rltopsp1 X0)\Rightarrow((l1\_rlvect\_1 X0)\wedge(l1\_pre\_topc X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((\neg v2\_struct\_0 X0)\wedge((v13\_algstr\_0 X0)\wedge((v2\_rlvect\_1 X0)\wedge((v3\_rlvect\_1 X0)\wedge((v4\_rlvect\_1 X0)\wedge((v5\_rlvect\_1 X0)\wedge((v6\_rlvect\_1 X0)\wedge((v7\_rlvect\_1 X0)\wedge((v8\_rlvect\_1 X0)\wedge(l1\_rlvect\_1 X0))))))))))\wedge((m1\_subset\_1 X1 (u1\_struct\_0 X0))\wedge(m1\_subset\_1 X2 (u1\_struct\_0 X0))))\Rightarrow(m1\_subset\_1 (k1\_rltopsp1 X0 X1 X2) (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(m1\_subset\_1 (k19\_euclid X0 X1) (u1\_struct\_0 (k15\_euclid np\_2))) \quad (14)$$

Assume the following.

$$\forall X0.(v7\_ordinal1\ X0)\Rightarrow((v5\_rltopsp1\ (k15\_euclid\ X0))\wedge (l1\_rltopsp1\ (k15\_euclid\ X0))) \quad (15)$$

Assume the following.

$$\forall X0.(m1\_subset\_1\ X0\ k1\_numbers)\Rightarrow(v1\_xreal\_0\ X0) \quad (16)$$

Assume the following.

$$\forall X0.(v6\_membered\ X0)\Rightarrow(\forall X1.(m1\_subset\_1\ X1\ X0)\Rightarrow (v7\_ordinal1\ X1)) \quad (17)$$

**Theorem 1**

$$r1\_tarski\ (k1\_rltopsp1\ (k15\_euclid\ np\_2)\ (k19\_euclid\ (k1\_real\_1\ np\_1)\ (k1\_real\_1\ np\_3))\ (k19\_euclid\ (k1\_real\_1\ np\_1)\ np\_3))\ (k4\_subset\_1\ (u1\_struct\_0\ (k15\_euclid\ np\_2))\ (k1\_rltopsp1\ (k15\_euclid\ np\_2)\ (k19\_euclid\ (k1\_real\_1\ np\_1)\ (k1\_real\_1\ np\_3))\ (k19\_euclid\ (k1\_real\_1\ np\_1)\ np\_3))\ (k1\_rltopsp1\ (k15\_euclid\ np\_2)\ (k19\_euclid\ (k1\_real\_1\ np\_1)\ np\_3))\ (k19\_euclid\ np\_1\ np\_3))$$