

l117_xreal_1 (TM-
MukfW91fTQkZi3RN4EHYUjU3ApwjfV3tG)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 (k3_xcmplx_0 X1 X0) X2) \Rightarrow ((r1_xxreal_0 \\ & X0 k6_numbers) \vee (r1_xxreal_0 X1 (k7_xcmplx_0 X2 X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 X0 k6_numbers = k6_numbers) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k3_xcmplx_0 X0 (k5_xcmplx_0 X1) = k7_xcmplx_0 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k7_xcmplx_0 (k5_xcmplx_0 X0) (k5_xcmplx_0 X1) = k7_xcmplx_0 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge ((\neg r1_xxreal_0 X1 k6_numbers) \wedge (r1_xxreal_0 (k3_xcmplx_0 X0 X1) k6_numbers)))) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 k6_numbers X2)) \Rightarrow (r1_xxreal_0 (k3_xcmplx_0 X0 X2) (k3_xcmplx_0 X1 X2)))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ (v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge (\neg r1_xxreal_0 \\ X2 X1) \wedge (r1_xxreal_0 (k3_xcmplx_0 X2 X0) (k3_xcmplx_0 X1 X0)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg \\ r1_xxreal_0 k6_numbers X0) \wedge (\neg r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 \\ (k5_xcmplx_0 X1) (k5_xcmplx_0 X0)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k5_xcmplx_0 (k5_xcmplx_0 X0) = X0) \quad (9)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k5_xcmplx_0 X0)) \wedge \\ (v1_xreal_0 (k5_xcmplx_0 X0))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow ((\\ (X0 \neq k6_numbers) \Rightarrow ((X1 = k5_xcmplx_0 X0) \Leftrightarrow (k3_xcmplx_0 X0 X1 = np_1))) \wedge \\ ((X0 = k6_numbers) \Rightarrow ((X1 = k5_xcmplx_0 X0) \Leftrightarrow (X1 = k6_numbers)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (12)$$

Theorem 1

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 \\ X0 X1) \Rightarrow ((r1_xxreal_0 X0 k6_numbers) \vee (r1_xxreal_0 (k5_xcmplx_0 \\ X1) (k5_xcmplx_0 X0)))))) \end{aligned}$$