

l11_jordan1d (TMW-
bLPpP5pwfaVSWxgqUKk5hKQgUiCTSaUh)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $k9_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_xreal_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & ((v2_xreal_0 \ np_2) \wedge (m2_subset_1 \ np_2 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_2 \ k5_numbers) \wedge (m1_subset_1 \ np_2 \ k1_numbers)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 \ X0 \ k1_numbers) \wedge (v1_xreal_0 \ X1)) \Rightarrow (k9_real_1 \ X0 \ X1 = k6_xcmplx_0 \ X0 \ X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 \ X0 \ k1_numbers) \wedge (v1_xreal_0 \ X1)) \Rightarrow (k8_real_1 \ X0 \ X1 = k3_xcmplx_0 \ X0 \ X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v7_ordinal1 \ X0) \wedge (v7_ordinal1 \ X1)) \Rightarrow (k7_nat_d \ X0 \ X1 = k1_xreal_0 \ X0 \ X1) \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 \ X0 \ k5_numbers) \wedge (v7_ordinal1 \ X1)) \Rightarrow (k4_nat_1 \ X0 \ X1 = k3_xcmplx_0 \ X0 \ X1) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((r1_xxreal_0 np_2 X0) \Rightarrow (r1_xxreal_0 k6_numbers (k9_real_1 (k8_real_1 np_2 X0) np_2))) \quad (7)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (v7_ordinal1 (k3_xcmplx_0 X0 X1)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k8_real_1 X0 X1) k1_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 k6_numbers (k6_xcmplx_0 X0 X1)) \Rightarrow (k1_xreal_0 X0 X1 = k6_xcmplx_0 X0 X1)) \wedge ((\neg r1_xxreal_0 k6_numbers (k6_xcmplx_0 X0 X1)) \Rightarrow (k1_xreal_0 X0 X1 = k6_numbers)))) \quad (11)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (13)$$

Assume the following.

$$\forall X0.(v6_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v7_ordinal1 X1)) \quad (14)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow ((r1_xxreal_0 np_2 X0) \Rightarrow (k9_real_1 (k4_nat_1 np_2 X0) np_2 = k7_nat_d (k4_nat_1 np_2 X0) np_2))$$