

l11_topreal2
(TML2LSHekJE4Ue7NTmFoNEcm4deoxiR1gb4)

October 27, 2020

Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \quad (1)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (2)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (3)$$

Theorem 1 $k18_euclid (k19_euclid np_1 np_1) = np_1$.