

l127_jordan (TMGHxKTfiCmiY- BqExwgL14msSEYKbBkpbm1)

October 27, 2020

Let $v2_sppol_1 : \iota \Rightarrow o$ be given. Let $k1_rltopsp1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid \\ (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ ((k17_euclid X0 = k17_euclid X1) \Leftrightarrow (v2_sppol_1 (k1_rltopsp1 (k15_euclid \\ np_2) X0 X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k1_real_1 X0) k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k19_euclid X0 X1) (u1_struct_0 (k15_euclid np_2))) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$v2_sppol_1 (k1_rltopsp1 (k15_euclid np_2) (k19_euclid np_1 \\ (k1_real_1 np_3)) (k19_euclid np_1 np_3))$$