

112_neckla_2 (TMK-
twije9vEmJKtNRpSEMQLYNzF95rW72ZA)

October 27, 2020

Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k4_necklace : \iota \Rightarrow \iota$ be given. Let $np_4 : \iota$ be given. Let $k2_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $np_2 : \iota$ be given. Let $np_3 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (u1_struct_0 (k4_necklace X0) = X0) \quad (1)$$

Assume the following.

$$np_4 = k2_enumset1 k6_numbers np_1 np_2 np_3 \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_4) \wedge (m2_subset_1 np_4 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_4 k5_numbers) \wedge (m1_subset_1 np_4 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (5)$$

Theorem 1

$$u1_struct_0 (k4_necklace np_4) = k2_enumset1 k6_numbers np_1 np_2 np_3$$