

l12_qmax_1

(TMR2QeWJgwVbexpE2EtzshgqkAxV7NgXauP6)

October 27, 2020

Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k12_prob_1 : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k1_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $c3_qmax_1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $c2_qmax_1 : \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((X2 = k1_tarski \\ (k4_tarski X0 X1)) \Rightarrow ((k9_xtuple_0 X2 = k1_tarski X0) \wedge (k10_xtuple_0 \\ X2 = k1_tarski X1))) \end{aligned} \tag{1}$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. \forall X3. ((m1_subset_1 X2 \\ (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1)))) \Rightarrow (r2_relset_1 X0 X1 X2 X2) \end{aligned} \tag{3}$$

Assume the following.

$$k6_numbers = k1_xboole_0 \tag{4}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow \\ (k6_domain_1 X0 X1 = k1_tarski X1) \end{aligned} \tag{5}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(\\ k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge \\ ((\neg v1_xboole_0 X1)\wedge((m1_subset_1 X2 X0)\wedge(m1_subset_1 X3 X1))))\Rightarrow \\ (k1_domain_1 X0 X1 X2 X3 = k4_tarSKI X2 X3) \quad (8)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\exists X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ X0))\wedge(\neg v1_xboole_0 X1)) \quad (9)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k1_tarSKI (k4_tarSKI X0 X1)) \quad (11)$$

Assume the following.

$$\forall X0.\neg v1_xboole_0 (k1_zfmisc_1 X0) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(\neg v1_xboole_0 X1))\Rightarrow \\ (\neg v1_xboole_0 (k2_zfmisc_1 X0 X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow \\ (m1_subset_1 (k6_domain_1 X0 X1) (k1_zfmisc_1 X0)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge \\ ((\neg v1_xboole_0 X1)\wedge((m1_subset_1 X2 X0)\wedge(m1_subset_1 X3 X1))))\Rightarrow \\ (m1_subset_1 (k1_domain_1 X0 X1 X2 X3) (k2_zfmisc_1 X0 X1)) \quad (15)$$

Assume the following.

$$(v1_funct_1 c2_qmax_1)\wedge((v1_funct_2 c2_qmax_1 k12_prob_1 \\ k1_numbers)\wedge(m1_subset_1 c2_qmax_1 (k1_zfmisc_1 (k2_zfmisc_1 \\ k12_prob_1 k1_numbers)))) \quad (16)$$

Assume the following.

$$c3_qmax_1 = c2_qmax_1 \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.(v1_xboole_0 X0) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_xboole_0 X2)) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (19)$$

Theorem 1

$$\begin{aligned} & r2_relset_1 k5_numbers k5_numbers (k1_relset_1 (k2_zfmisc_1 \\ & k5_numbers k5_numbers) (k6_domain_1 (k2_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers k5_numbers) (k1_zfmisc_1 (k2_zfmisc_1 k12_prob_1 \\ & k1_numbers)))) (k1_domain_1 (k2_zfmisc_1 k5_numbers k5_numbers) \\ & (k1_zfmisc_1 (k2_zfmisc_1 k12_prob_1 k1_numbers)) (k1_domain_1 \\ & k5_numbers k5_numbers k6_numbers k6_numbers) c3_qmax_1))) (\\ & k6_domain_1 (k2_zfmisc_1 k5_numbers k5_numbers) (k1_domain_1 \\ & k5_numbers k5_numbers k6_numbers k6_numbers)) \end{aligned}$$