

l12_xxreal_3 (TMN-
QjiBQhLp6KEgysKVvVoM478Md3ndDNme)

October 27, 2020

Let $k1_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_0 : \iota$ be given. Let $k1_xxreal_0 : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$k1_xxreal_3 \ k1_xxreal_0 \ k1_xxreal_0 = k1_xxreal_0 \quad (1)$$

Assume the following.

$$v1_xxreal_0 \ k2_xxreal_0 \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 \ X0) \Rightarrow (\forall X1.(v1_xxreal_0 \ X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 \ X2) \Rightarrow (((v1_xreal_0 \ X0) \wedge (v1_xreal_0 \ X1)) \Rightarrow ((X2 = \\ & \quad k1_xxreal_3 \ X0 \ X1) \Leftrightarrow (\exists X3.(v1_xcmplx_0 \ X3) \wedge (\exists X4. \\ & (v1_xcmplx_0 \ X4) \wedge ((X0 = X3) \wedge ((X1 = X4) \wedge (X2 = k2_xcmplx_0 \ X3 \ X4)))))) \wedge \\ & \quad (((X0 = k1_xxreal_0) \wedge (X1 \neq k2_xxreal_0)) \vee ((X1 = k1_xxreal_0) \wedge \\ & \quad (X0 \neq k2_xxreal_0))) \Rightarrow ((X2 = k1_xxreal_3 \ X0 \ X1) \Leftrightarrow (X2 = k1_xxreal_0))) \wedge \\ & \quad (((X0 = k2_xxreal_0) \wedge (X1 \neq k1_xxreal_0)) \vee ((X1 = k2_xxreal_0) \wedge \\ & \quad (X0 \neq k1_xxreal_0))) \Rightarrow ((X2 = k1_xxreal_3 \ X0 \ X1) \Leftrightarrow (X2 = k2_xxreal_0))) \wedge \\ & \quad (\neg(\neg(v1_xreal_0 \ X0) \wedge (v1_xreal_0 \ X1)) \wedge (\neg(X0 = k1_xxreal_0) \wedge \\ & \quad (X1 \neq k2_xxreal_0)) \wedge (\neg(X1 = k1_xxreal_0) \wedge (X0 \neq k2_xxreal_0)) \wedge \\ & \quad (\neg(X0 = k2_xxreal_0) \wedge (X1 \neq k1_xxreal_0)) \wedge (\neg(X1 = k2_xxreal_0) \wedge \\ & \quad (X0 \neq k1_xxreal_0))) \wedge (\neg(X2 = k1_xxreal_3 \ X0 \ X1) \Leftrightarrow (X2 = k6_numbers)))))) \end{aligned} \quad (3)$$

Assume the following.

$$k1_xxreal_0 = k1_numbers \quad (4)$$

Theorem 1 $k1_xxreal_3 \ k2_xxreal_0 \ k2_xxreal_0 = k2_xxreal_0$.