

l133_jordan
(TMQGKFK8S2zrD1ZgPwADQsy2nmMEjpUzLjR)

October 27, 2020

Let $v1_sppol_1 : \iota \Rightarrow o$ be given. Let $k1_rltopsp1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((k18_euclid X0 = k18_euclid X1) \Leftrightarrow (v1_sppol_1 (k1_rltopsp1 (k15_euclid \\ & \quad np_2) X0 X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (6)$$

Assume the following.

$$k18_euclid (k19_euclid\ np_1 (k1_real_1\ np_3)) = k1_real_1\ np_3 \quad (7)$$

Assume the following.

$$k18_euclid (k19_euclid\ k6_numbers (k1_real_1\ np_3)) = k1_real_1\ np_3 \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_numbers) \Rightarrow (m1_subset_1\ (k1_real_1\ X0)\ k1_numbers) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0\ X0) \wedge (v1_xreal_0\ X1)) \Rightarrow (m1_subset_1\ (k19_euclid\ X0\ X1)\ (u1_struct_0\ (k15_euclid\ np_2))) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_numbers) \Rightarrow (v1_xreal_0\ X0) \quad (11)$$

Theorem 1

$$v1_sppol_1 (k1_rltopsp1 (k15_euclid\ np_2) (k19_euclid\ np_1 (k1_real_1\ np_3)) (k19_euclid\ k6_numbers (k1_real_1\ np_3)))$$