

l133_toprealb (TMURT- mAJM4ogdVqSKKmxF6FTrv691FxBH2H)

October 27, 2020

Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k1_fcont_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $k10_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k32_sin_cos : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k31_sin_cos : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (k2_relset_1 k1_numbers (k1_fcont_1 X0 X1) = k1_numbers))) \quad (2)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (3)$$

Assume the following.

$$((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (4)$$

Assume the following.

$$\neg v1_xboole_0 np_2 \quad (5)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 \ np_1 \tag{7}$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 \ X0 \ k1_numbers)\wedge(v1_xreal_0 \ X1))\Rightarrow(k8_real_1 \ X0 \ X1 = k3_xcmplx_0 \ X0 \ X1) \tag{8}$$

Assume the following.

$$k6_numbers = k1_xboole_0 \tag{9}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{10}$$

Assume the following.

$$k32_sin_cos = k31_sin_cos \tag{11}$$

Assume the following.

$$\forall X0.(m1_subset_1 \ X0 \ k1_numbers)\Rightarrow(k1_real_1 \ X0 = k4_xcmplx_0 \ X0) \tag{12}$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 \ X0 \ k1_numbers)\wedge(v1_xreal_0 \ X1))\Rightarrow(k10_real_1 \ X0 \ X1 = k7_xcmplx_0 \ X0 \ X1) \tag{13}$$

Assume the following.

$$\forall X0.\forall X1.(((\neg v1_xboole_0 \ X0)\wedge(v1_xcmplx_0 \ X0))\wedge((\neg v1_xboole_0 \ X1)\wedge(v1_xcmplx_0 \ X1)))\Rightarrow(\neg v1_xboole_0 \ (k7_xcmplx_0 \ X0 \ X1)) \tag{14}$$

Assume the following.

$$\forall X0.\forall X1.(((\neg v1_xboole_0 \ X0)\wedge(v1_xcmplx_0 \ X0))\wedge((\neg v1_xboole_0 \ X1)\wedge(v1_xcmplx_0 \ X1)))\Rightarrow(\neg v1_xboole_0 \ (k3_xcmplx_0 \ X0 \ X1)) \tag{15}$$

Assume the following.

$$\forall X0.((\neg v1_xboole_0 \ X0)\wedge(v1_xcmplx_0 \ X0))\Rightarrow((\neg v1_xboole_0 \ (k4_xcmplx_0 \ X0))\wedge(v1_xcmplx_0 \ (k4_xcmplx_0 \ X0))) \tag{16}$$

Assume the following.

$$v1_xboole_0 \ k1_xboole_0 \tag{17}$$

Assume the following.

$$(v1_xreal_0 \ k31_sin_cos)\wedge(v2_xreal_0 \ k31_sin_cos) \tag{18}$$

Assume the following.

$$\forall X0.((\neg v3_xxreal_0 X0) \wedge (v1_xreal_0 X0)) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (\neg v2_xxreal_0 (k4_xcmplx_0 X0))) \quad (19)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k8_real_1 X0 X1) k1_numbers) \quad (20)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k1_real_1 X0) k1_numbers) \quad (21)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k10_real_1 X0 X1) k1_numbers) \quad (22)$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (23)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k5_numbers) \Rightarrow (\neg v3_xxreal_0 X0) \quad (24)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (25)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xcmplx_0 X0) \quad (26)$$

Theorem 1

$$k2_relset_1 k1_numbers (k1_fcont_1 (k1_real_1 (k10_real_1 np_1 (k8_real_1 np_2 k32_sin_cos))) np_1) = k1_numbers$$