

# l134\_jordan

(TMdFG5xBPhYR4hQrEZKP2Go5z31mW211Wg7)

October 27, 2020

Let  $v1\_sppol\_1 : \iota \Rightarrow o$  be given. Let  $k1\_rltopsp1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k19\_euclid : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_real\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $np\_3 : \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k17\_euclid : \iota \Rightarrow \iota$  be given. Let  $k18\_euclid : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow ((k17\_euclid (k19\_euclid X0 X1) = X0) \wedge (k18\_euclid (k19\_euclid X0 X1) = X1))) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 (k15\_euclid np\_2))) \Rightarrow \\ & (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 (k15\_euclid np\_2))) \Rightarrow \\ & ((k18\_euclid X0 = k18\_euclid X1) \Leftrightarrow (v1\_sppol\_1 (k1\_rltopsp1 (k15\_euclid \\ & \quad np\_2) X0 X1)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_3) \wedge (m2\_subset\_1 np\_3 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_3 k5\_numbers) \wedge (m1\_subset\_1 np\_3 k1\_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (6)$$

Assume the following.

$$\exists X0.(v1\_xboole\_0 X0) \wedge ((v1\_xcmplx\_0 X0) \wedge ((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X0))) \quad (7)$$

Assume the following.

$$k18\_euclid (k19\_euclid k6\_numbers np\_3) = np\_3 \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (m1\_subset\_1 (k1\_real\_1 X0) k1\_numbers) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (m1\_subset\_1 (k19\_euclid X0 X1) (u1\_struct\_0 (k15\_euclid np\_2))) \quad (10)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (11)$$

**Theorem 1**

$$v1\_sppol\_1 (k1\_rltopsp1 (k15\_euclid np\_2) (k19\_euclid (k1\_real\_1 np\_1) np\_3) (k19\_euclid k6\_numbers np\_3))$$