

l138_jordan

(TMavbT6ayPSzEjzAcxjHHhxBGCVE3XV7cHm)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rltopsp1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_3 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v2_sppol_1 : \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $np_0 : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X3.(m1_subset_1 X3 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (((v2_sppol_1 (k1_rltopsp1 (k15_euclid np_2) X0 X1)) \wedge ((v2_sppol_1 \\ & (k1_rltopsp1 (k15_euclid np_2) X2 X3)) \wedge ((k17_euclid X0 = k17_euclid \\ & X2) \wedge ((r1_xxreal_0 (k18_euclid X0) (k18_euclid X2)) \wedge ((r1_xxreal_0 \\ & (k18_euclid X2) (k18_euclid X3)) \wedge (r1_xxreal_0 (k18_euclid X3) \\ & (k18_euclid X1)))))) \Rightarrow (r1_tarski (k1_rltopsp1 (k15_euclid np_2) \\ & X2 X3) (k1_rltopsp1 (k15_euclid np_2) X0 X1)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$v1_xboole_0 \text{ np_}0 \quad (5)$$

Assume the following.

$$r1_xxreal_0 (k4_xcmplx_0 \text{ np_}3) \text{ np_}0 \quad (6)$$

Assume the following.

$$r1_xxreal_0 \text{ np_}3 \text{ np_}3 \quad (7)$$

Assume the following.

$$r1_xxreal_0 \text{ np_}0 \text{ np_}3 \quad (8)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (k1_real_1 X0 = k4_xcmplx_0 X0) \quad (10)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (11)$$

Assume the following.

$$k18_euclid (k19_euclid (k1_real_1 \text{ np_}1) k6_numbers) = k6_numbers \quad (12)$$

Assume the following.

$$k17_euclid (k19_euclid (k1_real_1 \text{ np_}1) k6_numbers) = k1_real_1 \text{ np_}1 \quad (13)$$

Assume the following.

$$v2_sppol_1 (k1_rltopsp1 (k15_euclid \text{ np_}2) (k19_euclid (k1_real_1 \text{ np_}1) k6_numbers) (k19_euclid (k1_real_1 \text{ np_}1) \text{ np_}3)) \quad (14)$$

Assume the following.

$$v2_sppol_1 (k1_rltopsp1 (k15_euclid \text{ np_}2) (k19_euclid (k1_real_1 \text{ np_}1) (k1_real_1 \text{ np_}3)) (k19_euclid (k1_real_1 \text{ np_}1) \text{ np_}3)) \quad (15)$$

Assume the following.

$$k18_euclid (k19_euclid (k1_real_1 \text{ np_}1) (k1_real_1 \text{ np_}3)) = k1_real_1 \text{ np_}3 \quad (16)$$

Assume the following.

$$k17_euclid (k19_euclid (k1_real_1 \text{ np_}1) (k1_real_1 \text{ np_}3)) = k1_real_1 \text{ np_}1 \quad (17)$$

Assume the following.

$$k18_euclid (k19_euclid (k1_real_1 np_1) np_3) = np_3 \quad (18)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (v1_xreal_0 (k4_xcmplx_0 X0))) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k19_euclid X0 X1) (u1_struct_0 (k15_euclid np_2))) \quad (20)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (21)$$

Theorem 1

$$r1_tarski (k1_rltopsp1 (k15_euclid np_2) (k19_euclid (k1_real_1 np_1) k6_numbers) (k19_euclid (k1_real_1 np_1) np_3)) (k1_rltopsp1 (k15_euclid np_2) (k19_euclid (k1_real_1 np_1) (k1_real_1 np_3)) (k19_euclid (k1_real_1 np_1) np_3))$$