

113_zfmodel2 (TMNkmCTpqDvYrwFPmhB- hXPu6vStEzeX7tii)

October 27, 2020

Let $k2_zf_lang : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $np_4 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k5_numbers) \Rightarrow ((k2_zf_lang X0 = k2_zf_lang X1) \Rightarrow (X0 = X1))) \quad (1)$$

Assume the following.

$$((v2_xxreal_0 np_4) \wedge (m2_subset_1 np_4 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_4 k5_numbers) \wedge (m1_subset_1 np_4 k1_numbers)) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \quad (3)$$

Theorem 1 $k2_zf_lang np_3 \neq k2_zf_lang np_4$.