

l141_quaterni

(TMJcfcB12XHnrkL1Y4nwMfmV4zqejUMm)

October 27, 2020

Let $v1_quaterni : \iota \Rightarrow o$ be given. Let $k29_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k28_quaterni : \iota \Rightarrow \iota$ be given. Let $k9_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_quaterni : \iota \Rightarrow \iota$ be given. Let $k26_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_quaterni : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_quaterni X0) \wedge (v1_quaterni X1)) \Rightarrow (k29_quaterni X0 X1 = k9_quaterni X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. (v1_quaterni X0) \Rightarrow (k28_quaterni X0 = k8_quaterni X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_quaterni X0) \wedge (v1_quaterni X1)) \Rightarrow (k26_quaterni X0 X1 = k7_quaterni X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. (v1_quaterni X0) \Rightarrow (\forall X1. (v1_quaterni X1) \Rightarrow (X0 = k26_quaterni (k29_quaterni X0 X1) X1)) \quad (4)$$

Assume the following.

$$\forall X0. (v1_quaterni X0) \Rightarrow (\forall X1. (v1_quaterni X1) \Rightarrow (X0 = k29_quaterni (k26_quaterni X0 X1) X1)) \quad (5)$$

Assume the following.

$$\forall X0. (v1_quaterni X0) \Rightarrow (k8_quaterni (k8_quaterni X0) = X0) \quad (6)$$

Assume the following.

$$\forall X0. (v1_quaterni X0) \Rightarrow (k28_quaterni (k28_quaterni X0) = X0) \quad (7)$$

Assume the following.

$$\forall X0.(v1_quaterni\ X0)\Rightarrow(v1_quaterni\ (k8_quaterni\ X0)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_quaterni\ X0)\wedge(v1_quaterni\ X1))\Rightarrow(m1_subset_1\ (k29_quaterni\ X0\ X1)\ k1_quaterni) \quad (9)$$

Assume the following.

$$\forall X0.(v1_quaterni\ X0)\Rightarrow(\forall X1.(v1_quaterni\ X1)\Rightarrow(k9_quaterni\ X0\ X1 = k7_quaterni\ X0\ (k8_quaterni\ X1))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_quaterni\ X0)\wedge(v1_quaterni\ X1))\Rightarrow(k26_quaterni\ X0\ X1 = k26_quaterni\ X1\ X0) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_quaterni)\Rightarrow(v1_quaterni\ X0) \quad (12)$$

Theorem 1

$$\forall X0.(v1_quaterni\ X0)\Rightarrow(\forall X1.(v1_quaterni\ X1)\Rightarrow(k29_quaterni\ X0\ X1 = k28_quaterni\ (k29_quaterni\ X1\ X0)))$$