

## l14\_comseq\_3

(TMbE4grQXEtD3B4k46s1VsC3WvtXep43ZCu)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k2\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k17\_complex1 : \iota \Rightarrow \iota$  be given. Let  $k8\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k55\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_valued\_0 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k54\_valued\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_membered : \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k17\_complex1 X0)) \quad (1)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_valued\_0 X0))) \Rightarrow (\forall X1.k1\_funct\_1 (k54\_valued\_1 X0) X1 = k17\_complex1 (k1\_funct\_1 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 k5\_numbers X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \wedge (v7\_ordinal1 X2)) \Rightarrow (k8\_nat\_1 X0 X1 X2 = k1\_funct\_1 X1 X2) \quad (4)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (5)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_membered X1)\wedge((v1\_funct\_1 X2)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))))\Rightarrow(k55\_valued\_1 X0 X1 X2 = k54\_valued\_1 X2) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((\neg v1\_xboole\_0 X1)\wedge(v1\_membered X1))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 X2 X0 X1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))))\Rightarrow((v1\_funct\_1 (k54\_valued\_1 X2))\wedge(v1\_partfun1 (k54\_valued\_1 X2) X0)) \quad (8)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_valued\_0 X0)))\Rightarrow(v1\_xcmplx\_0 (k1\_funct\_1 X0 X1)) \quad (10)$$

Assume the following.

$$\neg v1\_xboole\_0 k2\_numbers \quad (11)$$

Assume the following.

$$\neg v1\_xboole\_0 k1\_numbers \quad (12)$$

Assume the following.

$$v1\_membered k2\_numbers \quad (13)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_membered X1)\wedge((v1\_funct\_1 X2)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))))\Rightarrow((v1\_funct\_1 (k55\_valued\_1 X0 X1 X2))\wedge(m1\_subset\_1 (k55\_valued\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k1\_numbers)))) \quad (15)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v1\_partfun1 X2 X0) \Rightarrow (v1\_funct\_2 X2 X0 X1)) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.(v1\_membered X1) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_valued\_0 X2)) \quad (19)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k2\_numbers) \wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k2\_numbers)))))) \Rightarrow \\ & (\forall X1.(m2\_subset\_1 X1 k1\_numbers k5\_numbers) \Rightarrow ((k17\_complex1 \\ & (k8\_nat\_1 k2\_numbers X0 X1) = k8\_nat\_1 k1\_numbers (k55\_valued\_1 \\ & k5\_numbers k2\_numbers X0) X1) \wedge (r1\_xxreal\_0 k6\_numbers (k8\_nat\_1 \\ & k1\_numbers (k55\_valued\_1 k5\_numbers k2\_numbers X0) X1)))) \end{aligned}$$