

l14_pdiff_7 (TMXAKAxHkqH- CatwxXZu8oKFeQLXgKaNX1YG)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_finseq_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (\forall X1. \forall X2. (v7_ordinal1 X2) \Rightarrow (k3_finseq_1 (k2_funct_7 \\ X0 X2 X1) = k3_finseq_1 X0)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_2 X1 X0) \Rightarrow (\forall X2. (m2_finseq_2 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (k3_finseq_1 X0 = k1_card_1 X0) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. \forall X3. ((\neg v1_xboole_0 X0) \wedge \\ ((m1_finseq_1 X1 X0) \wedge ((v7_ordinal1 X2) \wedge (m1_subset_1 X3 X0)))) \Rightarrow \\ (k1_finseq_7 X0 X1 X2 X3 = k2_funct_7 X1 X2 X3) \end{aligned} \quad (6)$$

Assume the following.

$$v6_membered\ k4_ordinal1 \quad (7)$$

Assume the following.

$$\neg v1_xboole_0\ k1_numbers \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2\ X1\ X0)\Rightarrow(\forall X2.(m2_finseq_2\ X2\ X0\ X1)\Rightarrow(m2_finseq_1\ X2\ X0)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1\ X1\ X0)\Rightarrow((v1_relat_1\ X1)\wedge(v1_funct_1\ X1)\wedge(v1_finseq_1\ X1)) \quad (10)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(m1_finseq_2\ (k1_euclid\ X0)\ k1_numbers) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(v3_card_1\ X1\ X0)\Leftrightarrow(k1_card_1\ X1 = X0) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k1_euclid\ X0))\Rightarrow(v3_card_1\ X1\ X0)) \quad (13)$$

Assume the following.

$$\forall X0.(v6_membered\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v7_ordinal1\ X1)) \quad (14)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v1_xboole_0\ X0)\wedge(m1_subset_1\ X0\ k5_numbers))\Rightarrow \\ & (\forall X1.(m2_finseq_2\ X1\ k1_numbers\ (k1_euclid\ X0))\Rightarrow(\forall X2. \\ & (m1_subset_1\ X2\ k1_numbers)\Rightarrow(\forall X3.(m1_subset_1\ X3\ k5_numbers)\Rightarrow \\ & (k3_finseq_1\ (k1_finseq_7\ k1_numbers\ X1\ X3\ X2) = X0)))) \end{aligned}$$