

# l16\_cqc\_the2 (TMdPP- SEXWU85a5zSguQdjMBBMcXqwM9u8ti)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k3\_cqc\_lang : \iota \Rightarrow \iota$  be given. Let  $v2\_cqc\_the1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (k9\_cqc\_lang X0 X1 X2 = k8\_cqc\_lang X0 (k6\_cqc\_lang \\ X0 X1) X2))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (v2\_cqc\_the1 (k8\_cqc\_lang X0 (k9\_cqc\_lang \\ X0 X1 X2) (k8\_cqc\_lang X0 (k6\_cqc\_lang X0 X1) X2)) X0))) \end{aligned} \tag{2}$$

## Theorem 1

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (v2\_cqc\_the1 (k8\_cqc\_lang X0 (k8\_cqc\_lang \\ X0 (k6\_cqc\_lang X0 X1) X2) (k9\_cqc\_lang X0 X1 X2)) X0))) \end{aligned}$$