

l16_ens_1

(TMLNNWp53EdmU9MVyGRBRyETmDAv3wWAJrd)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_ens_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_ens_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k4_ens_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k2_ens_1 \\ & X0)) \Rightarrow ((\neg (k4_ens_1 X0 X1 = k1_xboole_0) \wedge (k3_ens_1 X0 X1 \neq k1_xboole_0)) \wedge \\ & ((v1_funct_1 (k2_xtuple_0 X1)) \wedge (v1_funct_2 (k2_xtuple_0 X1) \\ & (k3_ens_1 X0 X1) (k4_ens_1 X0 X1)) \wedge (m1_subset_1 (k2_xtuple_0 X1) \\ & (k1_zfmisc_1 (k2_zfmisc_1 (k3_ens_1 X0 X1) (k4_ens_1 X0 X1))))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (v1_xboole_0 (k9_xtuple_0 X0)) \quad (4)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow (((X1 \neq k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 \\ & X1) \Leftrightarrow (X0 = k1_relset_1 X0 X2))) \wedge ((X1 = k1_xboole_0) \Rightarrow ((v1_funct_2 \\ & X2 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X1)\Rightarrow((v5_relat_1 X1 X0)\Leftrightarrow(r1_tarski (k10_xtuple_0 X1) X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (9)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k2_ens_1 X0))\Rightarrow((k9_xtuple_0 (k2_xtuple_0 X1) = k3_ens_1 X0 X1)\wedge(r1_tarski (k10_xtuple_0 (k2_xtuple_0 X1)) (k4_ens_1 X0 X1))))$$