

l16_goboard5 (TMLeG- PYXz5QGUpP6Kz1znzArPUHWDCVwVTc)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_goboard1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge ((v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \quad (4)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (5)$$

Assume the following.

$$\forall X0. (m1_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow (m2_finseq_1 (k1_goboard1 X0) k1_numbers) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow ((X1 = k3_finseq_1 \\ & X0) \Leftrightarrow (k2_finseq_1 X1 = k9_xtuple_0 X0))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m2_finseq_1 X1 k1_numbers) \Rightarrow ((X1 = k1_goboard1 X0) \Leftrightarrow \\ & ((k3_finseq_1 X1 = k3_finseq_1 X0) \wedge (\forall X2.(m2_subset_1 X2 \\ & k1_numbers k5_numbers) \Rightarrow ((X2 \in k4_finseq_1 X1) \Rightarrow (k1_seq_1 X1 X2 = \\ & k17_euclid (k7_partfun1 (u1_struct_0 (k15_euclid np_2)) X0 X2))))))))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (k4_finseq_1 (k1_goboard1 X0) = k4_finseq_1 X0) \end{aligned}$$